



## Review

# The dual face of connexin-based astroglial $\text{Ca}^{2+}$ communication: A key player in brain physiology and a prime target in pathology<sup>☆</sup>



Marijke De Bock<sup>a,1</sup>, Elke Decrock<sup>a,\*,1,2</sup>, Nan Wang<sup>a</sup>, Mélissa Bol<sup>a</sup>, Mathieu Vinken<sup>b</sup>, Geert Bultynck<sup>c</sup>, Luc Leybaert<sup>a</sup>

<sup>a</sup> Department of Basic Medical Sciences, Physiology group, Faculty of Medicine and Health Sciences, Ghent University, B-9000 Ghent, Belgium

<sup>b</sup> Department of Toxicology, Center for Pharmaceutical Research, Faculty of Medicine and Pharmacy, Vrije Universiteit Brussel, B-1090 Brussels, Belgium

<sup>c</sup> Department of Cellular and Molecular Medicine, Laboratory of Molecular and Cellular Signalling, KU Leuven, Campus Gasthuisberg O/N-1 bus 802, B-3000 Leuven, Belgium

## ARTICLE INFO

### Article history:

Received 31 January 2014

Received in revised form 11 April 2014

Accepted 12 April 2014

Available online 21 April 2014

### Keywords:

$\text{Ca}^{2+}$

Connexin

Intercellular communication

Neuro-glio-vascular unit

Astrocyte

Neurodegeneration

## ABSTRACT

For decades, studies have been focusing on the neuronal abnormalities that accompany neurodegenerative disorders. Yet, glial cells are emerging as important players in numerous neurological diseases. Astrocytes, the main type of glia in the central nervous system, form extensive networks that physically and functionally connect neuronal synapses with cerebral blood vessels. Normal brain functioning strictly depends on highly specialized cellular cross-talk between these different partners to which  $\text{Ca}^{2+}$ , as a signaling ion, largely contributes. Altered intracellular  $\text{Ca}^{2+}$  levels are associated with neurodegenerative disorders and play a crucial role in the glial responses to injury. Intracellular  $\text{Ca}^{2+}$  increases in single astrocytes can be propagated toward neighboring cells as intercellular  $\text{Ca}^{2+}$  waves, thereby recruiting a larger group of cells. Intercellular  $\text{Ca}^{2+}$  wave propagation depends on two, parallel, connexin (Cx) channel-based mechanisms: *i*) the diffusion of inositol 1,4,5-trisphosphate through gap junction channels that directly connect the cytoplasm of neighboring cells, and *ii*) the release of paracrine messengers such as glutamate and ATP through hemichannels ('half of a gap junction channel'). This review gives an overview of the current knowledge on Cx-mediated  $\text{Ca}^{2+}$  communication among astrocytes as well as between astrocytes and other brain cell types in physiology and pathology, with a focus on the processes of neurodegeneration and reactive gliosis. Research on Cx-mediated astroglial  $\text{Ca}^{2+}$  communication may ultimately shed light on the development of targeted therapies for neurodegenerative disorders in which astrocytes participate. This article is part of a Special Issue entitled: Calcium signaling in health and disease. Guest Editors: Geert Bultynck, Jacques Haiech, Claus W. Heizmann, Joachim Krebs, and Marc Moreau.

© 2014 Elsevier B.V. All rights reserved.

## 1. Introduction

In the brain, multiple billions of cells communicate with one another in order to process highly complex information. While a major part of the intercellular communication occurs via neuronal synapses, glial

cells encompassing astrocytes, oligodendrocytes, tanycytes, radial glia and microglia, are critically involved in shaping cell-to-cell communication in the central nervous system (CNS) [1]. Cell–cell communication occurs between neurons, astrocytes, oligodendrocytes, activated microglia, vascular cells and ependymal cells [2] and is in large part

**Abbreviations:** A $\beta$ , amyloid- $\beta$ ; AD, Alzheimer's disease; ALS, amyotrophic lateral sclerosis; AQP4, aquaporin 4; APP, amyloid precursor protein; BAPTA, 1,2-bis(2-aminophenoxy)ethane-N,N,N',N'-tetraacetic acid; BBB, blood–brain barrier;  $[\text{Ca}^{2+}]_e$ , extracellular  $\text{Ca}^{2+}$  concentration;  $[\text{Ca}^{2+}]_i$ , cytoplasmic  $\text{Ca}^{2+}$  concentration; cADPR, cyclic adenosine diphosphoribose; CNS, central nervous system; CRACs,  $\text{Ca}^{2+}$  release activated channels; CSD, cortical spreading depression; Cx, connexin; EAE, experimental autoimmune encephalomyelitis; ER, endoplasmic reticulum; GFAP, glial fibrillary acidic protein; GJ, gap junction; GJIC, gap junction-mediated intercellular coupling; GPCR, G-protein-coupled receptor; HC, hemichannel; ILIC, inositol 1,4,5-trisphosphate-induced  $\text{Ca}^{2+}$  release; ICW, intercellular  $\text{Ca}^{2+}$  wave; IL-1 $\beta$ , interleukin-1 $\beta$ ; IP $_3$ , inositol 1,4,5-trisphosphate; IP $_3$ R, inositol 1,4,5-trisphosphate receptor;  $[\text{K}^+]_e$ , extracellular  $\text{K}^+$  concentration; KO, knock-out; L-AAA, L-alpha-aminoadipic acid; LPS, lipopolysaccharide; MAM, mitochondria-associated endoplasmic reticulum membrane; MCU, mitochondrial  $\text{Ca}^{2+}$  uniporter; GluR, glutamate receptor; MW, molecular weight; NAD $^+$ , nicotinamide adenine dinucleotide; NCX,  $\text{Na}^+/\text{Ca}^{2+}$  exchanger; NFAT, nuclear factor of activated T-cells; NFTs, neurofibrillary tangles; NGVU, neuro-glio-vascular unit; NMDAR, N-methyl-D-aspartate receptor; NO, nitric oxide; Panx, Pannexin; PLC, phospholipase C; PGE $_2$ , prostaglandin E $_2$ ; PS, presenilins; PDS, paroxysmal depolarization shift; PTP, permeability transition pore; ROS, reactive oxygen species; RyR, ryanodine receptor; SMC, smooth muscle cell; SOCE, store-operated  $\text{Ca}^{2+}$  entry; TNF- $\alpha$ , tumor necrosis factor- $\alpha$ ; TRP channel, transient receptor potential channel; VDAC, voltage-dependent anion channel

<sup>☆</sup> This article is part of a Special Issue entitled: Calcium signaling in health and disease. Guest Editors: Geert Bultynck, Jacques Haiech, Claus W. Heizmann, Joachim Krebs, and Marc Moreau.

\* Corresponding author at: Dept. Basic Medical Sciences, Physiology group, Ghent University, De Pintelaan 185 (Block B, 3rd floor), B-9000 Ghent, Belgium. Tel.: +32 9 332 39 73; fax: +32 9 332 30 59.

E-mail address: [Elke.Decrock@gmail.com](mailto:Elke.Decrock@gmail.com) (E. Decrock).

<sup>1</sup> Contributed equally, share first authorship.

<sup>2</sup> E.D. is a postdoctoral research fellow of the Fund for Scientific Research Flanders (FWO-Vlaanderen), Belgium.

mediated by paracrine signaling, involving the release, diffusion and interaction of a messenger molecule with its receptor. Additionally, more direct signaling is possible by the exchange of ions and low molecular weight (MW) molecules (<1.5 kDa) via gap junctions (GJs) that cluster in plaques and directly connect the cytoplasm of neighboring cells. GJs are formed by the docking of two hemichannels (HCs). The plasma membrane normally also contains HCs *not* incorporated into GJs which are sometimes referred to as unapposed HCs. These HCs are commonly closed but may be activated in response to various stimuli, thereby opening a pore that allows ionic and molecular exchange between the intra- and extracellular environment. Opening of unapposed HCs is a potential pathway for the diffusive release of messenger molecules like ATP and glutamate and may therefore contribute to paracrine signaling [3]. GJs and HCs are composed of connexin (Cx) subunits arranged in a, respectively dodecameric or hexameric, configuration. Cxs are tetraspan membrane proteins of which 21 human species have been identified and named according to their MW. Pannexins (Panxs) are another family of channel-forming proteins that have topological similarity to Cx hemichannels. Currently, three human Panxs (Panx1–3) have been identified. Both Cx43 and Panx1 are predominant in the CNS (Table 1) [4–8].

Normal brain functioning strictly depends on highly specialized cellular cross-talk to which calcium ions ( $\text{Ca}^{2+}$ ) largely contribute as a second messenger. In the brain, Cx/Panx channels crucially contribute to the cell-to-cell propagation of increases in the cytoplasmic free  $\text{Ca}^{2+}$  concentration ( $[\text{Ca}^{2+}]_i$ ), known as intercellular  $\text{Ca}^{2+}$  waves (ICWs) [9–11]. Neuronal communication primarily relies on action potential firing whereby  $[\text{Ca}^{2+}]_i$  changes have a critical role in regulating neurotransmitter release, synaptic plasticity and gene expression [12–14]. For electrically non-excitable cells such as glia and vascular endothelial cells,  $\text{Ca}^{2+}$  signals are the major fast (on a second time scale) system available for intra- as well as intercellular signaling. ICWs typically emerge in glial networks [11,15] as has been demonstrated in vivo [16–19], bringing up the question whether these ICWs play an active role in normal brain functioning and information processing, particularly at the level of synaptic signaling and regulation of blood supply. In addition, a wide diversity of pathologies (e.g. brain ischemia [20], Alzheimer's disease (AD) [21,22], CNS traumatic injury [23,24] and epilepsy [25,26]) has been associated with abnormal intercellular  $\text{Ca}^{2+}$  signaling, which is linked to inflammatory conditions and unwarranted cell death, supporting a pathological function for ICWs. In this review, we discuss the currently available evidence that describes  $\text{Ca}^{2+}$  signaling in the brain, focusing on intra- and intercellular glial  $\text{Ca}^{2+}$  signaling and the contribution of Cx channels, both in physiology and pathology.

## 2. The neuro-glio-vascular unit

The term 'neurovascular unit' originally emerged as a concept to widen the scope of possibilities for therapeutic interventions in stroke (Report of the Stroke Progress Research Group, 2002, [www.ninds.nih.gov](http://www.ninds.nih.gov)) and has been instrumental in scaling down the classical neurocentric view in favor of a more integrated approach by including vascular and blood targets. We indulge in denoting this unit as the 'neuro-glio-

vascular unit' (NGVU) to underscore the importance of glial cells as central players linking neurons to the cerebral vasculature. The NGVU is composed of neurons, astrocytes, microglial cells, endothelial cells, pericytes, smooth muscle cells (SMCs) and circulating blood cells [27] (Fig. 1). Functionally, these cells interact to control synaptic communication, blood–brain barrier (BBB) function, local blood supply, neuronal development and surveillance/immune function [28].

Astrocytes are centrally positioned within the NGVU, where they form extensive networks that physically and functionally connect neuronal synapses with cerebral vessels [2]. At one end, astrocytes send their processes to neuronal synapses, where they join pre- and postsynaptic structures, altogether forming a 'tripartite synapse'. A single rodent astrocyte may contact an estimated 140,000 synapses while in the human brain, astrocytes are hypothesized to support and modulate the function of roughly two million synapses [29,30]. By surrounding the synaptic cleft, astrocytes sense neuronal signaling activities and modulate them by secreting gliotransmitters like ATP, glutamate, D-serine and adenosine [31–36]. At the other end, astrocytes project 'endfeet' onto brain capillary endothelial cells and pericytes, thereby enveloping 98% of the vascular surface area. At the synaptic as well as vascular contact points, astrocytes are additionally equipped with a large repertoire of neurotransmitter receptors [37] that sense synaptic, neurometabolic and vascular activities. The dual contact with neurons and cerebral vessels thus provides astrocytes with the unique architectural advantage for sensing neuronal activity, integrating electrochemical information and exchanging this information with the vasculature, for example, by releasing vasoactive substances such as nitric oxide (NO), prostacyclins, epoxyeicosatrienoic acids, glutamate, adenosine and ATP [38–40]. As such, the NGVU performs a homeostatic control over the neuronal milieu, which is required for proper functioning of neurons and their synapses.

## 3. Connexin signaling and glial network communication

Twelve Cxs and two Panxs have been identified in the various CNS cell types (Table 1). Each cell type expresses a unique combination of Cx species that typically depends on the brain region and on the developmental stage, with Cx43 being most ubiquitously expressed [6–8,41–43]. It is predominant in astrocytes, activated microglia, developing neurons, SMCs, endothelial cells and pericytes. Cx-mediated intercellular communication, being a typical feature of glial cells, is thought to play an important role in signal transmission at the NGVU, primarily via the formation of glial networks [44].


Astroglial cells prominently express Cxs and through extensive gap junctional coupling they establish astrocytic networks that support long-range communication and transport [44]. Importantly, interastroglial coupling is not uniform but endowed with anatomical or functional specificity, and is subject to modulation [45,46]. Cx43 and Cx30 are the main Cx species expressed in astrocytes and both contribute to GJ-mediated intercellular coupling (GJIC) between these cells [47–52]. Cx43 and Cx30 are primarily expressed in astrocytic processes that surround the synaptic cleft, but are also enriched in astrocytic endfeet. As GJs, these Cxs support intercellular trafficking of ions and small signaling molecules through perivascular and perisynaptic astroglial networks. Recent data obtained from transgenic mice with a targeted deletion of Cx26 in neuronal and glial cells, provided unequivocal evidence for the additional presence of Cx26 in astrocytic GJ plaques as well as for heterotypic plaques formed between astrocytes and oligodendrocytes [53]. Other Cxs (Cx40 and Cx45) have been detected in astrocytes, but are expressed at low levels [54,55]. The pattern of Cx expression in astrocytes is subject to alterations; e.g., Cx30 and Cx43 are upregulated in the presence of synaptically active neurons [51,56] and astrocyte-specific Cx43 knock-out (KO) is associated with a compensatory increase in Cx30 expression [57]. Under pathological conditions, Cx

**Table 1**  
Main connexin and pannexin species expressed in the neuroglivascular unit.

	Connexin	Pannexin
Astrocyte	Cx26, Cx30, Cx43	Panx1, Panx2 <sup>a</sup>
Microglia	Cx32, Cx36, Cx43 <sup>b</sup>	Panx1
Oligodendrocyte	Cx29, Cx32, Cx47	Panx1, Panx2
Neuron	Cx30.2, Cx31.1, Cx36, Cx40, Cx45	Panx1, Panx2
Endothelial cell	Cx37, Cx40, Cx43	ND
Pericyte	Cx37, Cx40, Cx43	ND

<sup>a</sup> De novo expression in reactive hippocampal astrocytes subjected to ischemia/reperfusion injury.

<sup>b</sup> Expressed in activated microglia, not in resting microglia.



**Fig. 1.** Schematic diagram of the neurogliovascular unit. Astrocytes send projections towards the synapse, where they form the tripartite synapse together with pre- and postsynaptic nerve terminals. At the other end, astrocyte endfeet enwrap the vascular wall from the capillary to the arteriolar level. In this way, neurons can influence the endothelium, using the astrocytes as intermediate players.

expression may additionally experience marked changes (see Section 5).

Several global and conditional Cx KO mice for the glial Cxs (Cx43, Cx30 and Cx26) are available and, as a consequence of the previously mentioned compensatory changes in the expression levels of other Cxs, KO mice deficient for multiple glial Cxs have been generated as well [45]. Targeted astrocytic ablation of Cx43 on a global Cx30 KO background renders mice that are largely devoid of astrocytic GJIC. The diffusion of glucose metabolites through the astrocytic network was abolished in hippocampal slices from these double KO mice [49]. Astrocytic networks exert homeostatic control over the composition of the extracellular environment, thereby influencing neuronal function and survival. Astrocytic GJIC is crucial for the supply and delivery of energy molecules and for the removal and spatial buffering of extracellular  $K^+$  and glutamate [2,52]. Increased neuronal activity augments astrocytic GJIC [49,58] and induces vasodilation of upstream pial arterioles. While in most perfusion beds, upstream signaling from capillaries to arterioles is mediated by the communication of electrical signals via endothelial GJs, there seems to be an alternative pathway in the brain consisting of the propagation of  $Ca^{2+}$  signals over astrocytic endfeet that strongly express Cx43 [59]. Work with Cx30/Cx43 double KO animals has additionally provided evidence that these Cxs are necessary for normal BBB integrity and function [48]. The evidence above strongly suggests that energy delivery, clearing and signaling functions of astroglial Cx channels are adapted to the level of neuronal activity. At the neuronal and vascular contact points, transport and signaling interactions with the glial network are mainly governed by mechanisms

other than GJs [50,60] although direct signaling through neuron–glial or endothelial–glial GJs has been reported in cell culture systems [61–64]. By means of immunolabeling techniques or functional assays, GJs between astrocytes and neurons have been detected in brain slices [65–67]. GJs between astrocytes and endothelial cells have been described in cell culture models [68] but they are normally not present at the healthy astrocyte–endothelial interface, presumably because of the presence of a continuous basal membrane. However, this does not exclude the possibility that heterocellular astrocyte–endothelial GJs can form under pathological conditions.

The participation of GJs in brain functioning has been abundantly reviewed but evidence for contribution of astrocytic HCs has only been accumulating over the past 10 years [69,70]. Before being incorporated into GJs, HCs form a bidirectional diffusion pathway for ions, small signaling molecules and energy substrates [3,69,71]. In this respect, Cx43 HCs have been demonstrated, mostly under *in vitro* conditions, to function as a release pathway for molecules such as ATP, glutamate, prostaglandin  $E_2$  ( $PGE_2$ ), nicotinamide adenine dinucleotide ( $NAD^+$ ) and inositol 1,4,5-trisphosphate ( $IP_3$ ) [72–77]. Cx43 HC opening has also been proposed to act as a pathway for the uptake of energy molecules in astrocytes in response to pro-inflammatory cytokines [78]. However, the biological significance of HCs is still a matter of debate as it is hard to understand how a poorly selective plasma membrane channel, with a conductance about twice the conductance of the corresponding GJ channel [79,80], can contribute to the normal functioning of a cell [81]. It was therefore hypothesized that HCs remain closed under most conditions to prevent leakage of metabolites and nutrients

and loss or entry of ions that would otherwise lead to cell dysfunction and cell death. Under resting conditions, Cx43 HCs exhibit a low open probability in cultured cells [79,82] that can be increased by exposure to various stimuli such as mechanical stimulation, changes in phosphorylation status [83], ischemic conditions [84], pro-inflammatory cytokines [78], oxidative stress [85], and an increase in  $[Ca^{2+}]_i$  [86] or decrease in extracellular  $Ca^{2+}$  concentration ( $[Ca^{2+}]_e$ ) [87]. Although most of these stimuli are associated with cellular stress, HC function has been associated with various aspects of the cellular life cycle, including cell proliferation, as well as with the normal functioning of various cell types [88]. Their involvement under normal non-pathological conditions is now supported by the observation that  $[Ca^{2+}]_i$  changes, in the physiological (submicromolar) range, trigger HC opening [86,89]. HC activity has also been recorded in cells exposed to endogenous ligands associated with cell proliferation and differentiation [83,90,91]. Nerve growth factor stimulated neurite outgrowth in PC12 cells was for instance mediated by HCs through ATP release [90]. Recently, Stehberg and co-workers pinpointed a contribution of astroglial Cx43 HCs in the basolateral amygdala to the consolidation of fear memory [92], suggesting a role of these channels in gliotransmitter release. Of note, functional Panx1 HCs have also been reported in astrocytes, but their role as gliotransmitter release channels has been less characterized and they do not form GJs between cultured astrocytes [93–95]. Hence, these channels are further referred to as Panx channels instead of Panx HCs [96].

Besides forming channels, Cxs also exert non-channel signaling functions [97,98]. Growing evidence suggests that Cx proteins directly control gene expression. Cx gene transfection is indeed known to affect the expression of genes encoding for growth-, differentiation- and cell death-modulating proteins [98]. The up- or down-regulation of various genes with diverse functions in the brain of Cx43-null mice indicate that Cxs regulate multiple cellular processes [99,100]. Irrespective of their channel-forming capacity, Cxs modulate P2Y receptor signaling [101, 102] and cytoskeletal proteins in astrocytes [103]. Recently, Cxs were found to be crucial for glial-guided neuronal migration without acting in the classical channel-mediated manner [104,105]. Instead, Cx26 and Cx43 GJs, expressed at the contact points between radial fibers and migrating neurons, provided dynamic adhesive contacts that interact with the internal cytoskeleton to enable leading process stabilization along radial fibers as well as the subsequent translocation of the nucleus [104].

#### 4. $Ca^{2+}$ signaling in the healthy brain

##### 4.1. General concepts of intra- and intercellular $Ca^{2+}$ signaling

$[Ca^{2+}]_i$  is regulated through a strongly organized interplay between  $Ca^{2+}$  entry from the extracellular space,  $Ca^{2+}$  release from intracellular storage sites, and  $Ca^{2+}$  extrusion out of the cells or its sequestration back into the stores. Most cytoplasmic  $Ca^{2+}$  is furthermore bound by  $Ca^{2+}$  buffering proteins such as parvalbumin, calretinin and calbindin [106]. As a result,  $[Ca^{2+}]_i$  in resting brain cells varies between 80 nM and 100 nM [21,107]. At high cytosolic levels,  $Ca^{2+}$  ions have a deleterious action; they cause aggregation of proteins and nucleic acids, affect the integrity of lipid membranes and initiate the precipitation of phosphates, and are thus incompatible with life [108]. Therefore, all living cells keep  $[Ca^{2+}]_i$  at low nanomolar levels, creating a steep concentration gradient between the extracellular environment (~1 mM [109–111]), the cytoplasm and the  $Ca^{2+}$  store content. Free cytoplasmic  $Ca^{2+}$  represents only a small part of the total cellular  $Ca^{2+}$  as most of the intracellular  $Ca^{2+}$  is sequestered in intracellular stores to prevent the toxic effects of a high  $[Ca^{2+}]_i$ . The endoplasmic reticulum (ER) is considered the predominant  $Ca^{2+}$  store as it contains approximately 70% of the cell's total  $Ca^{2+}$  reserve, corresponding to  $Ca^{2+}$  concentrations up to 0.5 mM [112–114]. Additionally, mitochondria play an important role in the  $Ca^{2+}$  household of cells [115]. Transport across the plasma membrane and membranes of the intracellular compartments

is governed by a family of  $Ca^{2+}$  channels and transporters that are based on the diffusion of  $Ca^{2+}$  down its concentration gradient (passive downhill), causing an increase in  $[Ca^{2+}]_i$ . The clearance of cytosolic  $Ca^{2+}$  depends on different energy-dependent transport mechanisms that act against the concentration gradient (active uphill) (Table 2) [116].

Variations in  $[Ca^{2+}]_i$  control a multitude of biological processes over a wide dynamic range including cell proliferation, differentiation, neurotransmitter release, secretion, synaptic plasticity, gene expression, immune responses, muscle contraction, endothelial permeability, apoptosis and many others [106]. To ensure specificity,  $Ca^{2+}$  signals are highly organized in time and space [10]. Rapid and highly localized  $Ca^{2+}$  spikes may co-occur or evolve into slower global  $[Ca^{2+}]_i$  changes that are either transient, repetitive or sustained. Neuronal  $[Ca^{2+}]_i$  rises mainly rely on  $Ca^{2+}$  entry via voltage- and receptor-operated plasma membrane channels, evoking highly localized and precisely timed  $[Ca^{2+}]_i$  increases that ensure a fast, focal and timed neurotransmitter release from synaptic terminals [12]. In contrast, macroglial (astrocytes and oligodendrocytes) as well as microglial  $Ca^{2+}$  signaling mechanisms mostly rely on  $Ca^{2+}$  release from the ER. This creates a longer-lasting  $[Ca^{2+}]_i$  elevation that is hypothesized to control a slower, and perhaps more sustained release of gliotransmitters [15]. The activation of store-operated  $Ca^{2+}$  entry (SOCE) following the depletion of the ER  $Ca^{2+}$  store, is equally important for shaping glial  $Ca^{2+}$  signals [117]. It gives rise to a sustained  $Ca^{2+}$  signal which may significantly outlast the periods of ER  $Ca^{2+}$  release induced by metabotropic receptor stimulation [17]. The key molecular players of SOCE are the stromal interaction molecules, which sense the luminal ER  $Ca^{2+}$  levels and control the opening of plasma membrane Orai channels [118,119]. In addition, Transient Receptor Potential (TRP) channels of the canonical family (TRPC) may contribute to SOCE [120], but this remains controversial [121].

The ER is the largest and most controlled intracellular  $Ca^{2+}$  store [122]. Generally,  $Ca^{2+}$  release from the ER is mediated by two families of receptor channels,  $IP_3$  receptors ( $IP_3$ Rs) and ryanodine receptors (RyRs), which are both represented by three isoforms ( $IP_3R1$ –3 and  $RyR1$ –3) [123–126].  $Ca^{2+}$  itself is an activator of RyR channels and a potent regulator of  $IP_3$ -induced  $Ca^{2+}$  release;  $Ca^{2+}$  thus provides a regenerative autocatalytic mechanism which is referred to as  $Ca^{2+}$ -induced  $Ca^{2+}$  release [127]. The primary activation mechanism of  $IP_3$ Rs is  $IP_3$  binding.  $IP_3$  is generated by phospholipase C (PLC), which is linked to a multitude of plasmalemmal metabotropic receptors. There are 13 different PLC isoforms, which are divided into six different classes ( $PLC\beta$ ,  $PLC\gamma$ ,  $PLC\delta$ ,  $PLC\epsilon$ ,  $PLC\zeta$  and  $PLC\eta$ ) based on their structure. These PLC proteins can be activated via different mechanisms including the stimulation of G-protein-coupled receptors (GPCRs), receptor and non-receptor tyrosine kinases, the activation of the small G-protein Ras, or by an increase in  $[Ca^{2+}]_i$  [128]. PLC catalyzes the cleavage of phosphoinositol-4,5-bisphosphate into diacylglycerol and  $IP_3$ . The latter diffuses inside the cell and binds its receptors located on the ER surface, resulting in  $IP_3$ -induced  $Ca^{2+}$  release (IICR) from the ER [116]. Astrocytes are equipped with a large repertoire of neurotransmitter receptors, many of which couple to the activation of PLC, production of  $IP_3$  and subsequent release of  $Ca^{2+}$  from internal stores [37,129]. At low levels of stimulation, single  $IP_3$ R channels open resulting in brief and localized  $Ca^{2+}$  blips, but usually, coordinated opening of clusters of  $IP_3$ Rs takes place, giving rise to  $Ca^{2+}$  puffs. Under certain conditions, these localized events may expand and recruit nearby ER  $Ca^{2+}$  channels to generate an intracellular  $Ca^{2+}$  wave sweeping over the cytoplasm thereby globalizing the  $Ca^{2+}$  response within the cell [130]. The spatio-temporal organization of  $Ca^{2+}$  signals thus tightly depends on the rearrangement of the  $Ca^{2+}$  channels on the ER surface, as well as their close contact with other  $Ca^{2+}$ -sequestering organelles such as mitochondria.

It becomes increasingly recognized that mitochondria play an important role in shaping the  $Ca^{2+}$  signals [115,131]. Consequent to proton extrusion by the electron transfer chain, required for ATP synthesis, mitochondria have an internally negative electrical potential



**Table 2**Mechanisms of  $\text{Ca}^{2+}$  transport between cytoplasm and extracellular environment or intracellular stores.

Passive downhill	Transport mode	Remarks
EC to cytoplasm	–Voltage-operated channel –Receptor-operated channel –Store-operated channel –Second messenger-operated channel –Non-selective channel	
ER to cytoplasm	–IP <sub>3</sub> R –RyR –ER leak channels <sup>a</sup>	
between cytoplasm and mitochondria	–MCU –VDAC –PTP	MCU is a highly selective $\text{Ca}^{2+}$ channel across the IMM while VDAC is the major regulator across the OMM PTP opening is associated with pathological conditions and results in $\text{Ca}^{2+}$ transport from mitochondria to the cytoplasm
Golgi to cytoplasm lysosomes and endosomes to cytoplasm	IP <sub>3</sub> R TPC	
Active uphill	Transport mode	Remarks
cytoplasm to EC cytoplasm to ER cytoplasm to Golgi	PMCA SERCA –SPCA –SERCA	
Secondary active uphill	Transport mode	Remarks
plasma membrane	NCX	The energy of the $\text{Na}^{+}$ or $\text{H}^{+}$ gradient is used to extrude $\text{Ca}^{2+}$ from the cell or from the mitochondria, however, it can work in reverse mode depending on the transmembrane $\text{Na}^{+}$ , $\text{H}^{+}$ or $\text{Ca}^{2+}$ gradients and membrane depolarization
mitochondrial membrane	–NCX – $\text{H}^{+}/\text{Ca}^{2+}$ antiporter	

This list summarizes the most important  $\text{Ca}^{2+}$ -permeable channels and pumps responsible for the  $\text{Ca}^{2+}$  transport between the compartments. Additional less-known mechanisms are reported to contribute but are not mentioned in the scheme.


(EC, extracellular environment; IMM, inner mitochondrial membrane; MCU, mitochondrial  $\text{Ca}^{2+}$  uniporter; NCX,  $\text{Na}^{+}/\text{Ca}^{2+}$  exchanger; OMM, outer mitochondrial membrane; PMCA, plasma membrane  $\text{Ca}^{2+}$  ATPase; PTP, permeability transition pore; SERCA, sarco-endoplasmic reticulum  $\text{Ca}^{2+}$  ATPase; SPCA, secretory pathway  $\text{Ca}^{2+}$  ATPase; TPC, two-pore channels; VDAC, voltage-dependent anion channel).

<sup>a</sup> ER leak channels remain unidentified up to date, but IP<sub>3</sub>R, RyR, TRP and Panx channels have been suggested to contribute to the  $\text{Ca}^{2+}$  leak.

that attracts  $\text{Ca}^{2+}$  to enter the mitochondrial matrix. Mitochondrial  $\text{Ca}^{2+}$  uptake is mediated by the voltage-dependent anion channel (VDAC) located at the outer mitochondrial membrane and the mitochondrial  $\text{Ca}^{2+}$  uniporter (MCU) which is situated in the inner mitochondrial membrane [132–136]. Mitochondria accumulate  $\text{Ca}^{2+}$  more effectively when they are close to sites of high  $\text{Ca}^{2+}$  concentration [137,138]. They are physically and functionally connected to the ER, separated by a distance in the range of 10–30 nm [139,140] that facilitates the uptake of  $\text{Ca}^{2+}$  released by the ER [141]. Contact sites between the two organelles are known as mitochondria-associated ER membranes or MAMs [142,143]. These are intracellular lipid raft-like structures intimately involved in cholesterol and phospholipid metabolism, and in  $\text{Ca}^{2+}$  homeostasis by providing direct transfer of ER-derived  $\text{Ca}^{2+}$  to the mitochondria [144–149]. Approximately 75 proteins have been reported to be associated with the MAMs [150]. Mitofusin 2 proteins are important for the tethering of the two organelles [151,152] and other proteins such as phosphofurin acidic cluster sorting protein 2, are reported to stabilize and regulate the interaction [153]. An interaction between the IP<sub>3</sub>R1 and VDAC1 via the chaperone ‘glucose regulated protein 75’ favors  $\text{Ca}^{2+}$  transfer [151]. Upon IICR, local microdomains with  $\text{Ca}^{2+}$  concentrations exceeding ~10  $\mu\text{M}$  (‘hotspots’) are created in the vicinity of mitochondria [137,141], favoring mitochondrial  $\text{Ca}^{2+}$  entry via the MCU that has an intrinsic low  $\text{Ca}^{2+}$  affinity ( $\text{EC}_{50}$  is 10–20  $\mu\text{M}$ ) [138,154]. A major role of mitochondrial  $\text{Ca}^{2+}$  entry and consequent increase of matrix  $[\text{Ca}^{2+}]$  consists in the stimulation of ATP production by activating  $\text{Ca}^{2+}$ -sensitive dehydrogenases of the Krebs cycle, regulating the activity of the ATP synthase and promoting the supply of oxidizable substrates [131,155–157]. Once the cytosolic  $\text{Ca}^{2+}$  has returned to its resting level, mitochondrial  $\text{Na}^{+}/\text{Ca}^{2+}$  exchangers (NCX) or  $\text{H}^{+}/\text{Ca}^{2+}$  antiporters transport  $\text{Ca}^{2+}$  from the mitochondrial matrix into the cytoplasm, from where it is either recycled to the ER or removed from the cell [158,159]. MAMs thus emerge as signaling hubs controlling oxidative cell energy metabolism and alterations at the MAMs are implicated

in a wide spectrum of diseases, including ischemia and neurodegenerative disorders (see Section 5, and [148,160]).

$\text{Ca}^{2+}$  waves are not always restricted to the cytosol of a single cell, but can cross the cell boundaries to form an ICW (Fig. 2). GJs offer the most direct route for ICW propagation by allowing the transfer of  $\text{Ca}^{2+}$  and IP<sub>3</sub>.  $\text{Ca}^{2+}$  diffusion in the cytoplasm is slow (diffusion constant in the order of 13 to 65  $\mu\text{m}^2/\text{s}$ ) due to the presence of slowly mobile  $\text{Ca}^{2+}$  binding proteins [161,162]. In contrast, the diffusion of IP<sub>3</sub> (MW 420 Da) is faster (diffusion constant ~283  $\mu\text{m}^2/\text{s}$ ) and GJs are more permeable to IP<sub>3</sub> than to  $\text{Ca}^{2+}$  (100-fold difference) [161,163]. Thus,  $\text{Ca}^{2+}$  can pass through GJs, but it reaches the adjacent cell with some delay compared to IP<sub>3</sub>. IP<sub>3</sub> therefore stands out as the primary  $\text{Ca}^{2+}$  signaling molecule passing via GJs [164]. Notably, GJs display selective permeability toward IP<sub>3</sub> depending on their Cx composition. Niessen and colleagues demonstrated that microinjection of IP<sub>3</sub> into HeLaCx32 cells triggered ICWs that were 3- to 4-fold more widespread than in HeLaCx26 cells and 2.5-fold larger than in HeLaCx43 cells [165]. Cyclic adenosine diphosphoribose (cADPR; MW 541 Da) is another GJ-permeable intercellular signaling molecule that triggers ICWs upon injection in astrocytes, but  $\text{Ca}^{2+}$  wave propagation is slower as compared to IP<sub>3</sub>-based ICWs [166]. cADPR is synthesized from NAD<sup>+</sup> by the ADP-ribosyl cyclase CD38, which is located on the cell surface and intracellular membranes. cADPR functions as a  $\text{Ca}^{2+}$  messenger by modulating the  $\text{Ca}^{2+}$  sensitivity of RyRs and appears to play an important role in  $\text{Ca}^{2+}$  signaling in the CNS, including in glial cells [167,168]. Apart from GJIC, non-junctional paracrine signaling also contributes to the propagation of ICWs (Fig. 2) [9–11]. The extracellular messengers involved are in most cases ATP or glutamate [169–173], but others such as NO [174] and NAD<sup>+</sup> [175] have also been suggested. Release of ATP and glutamate has been documented to occur via exocytosis or diffusion through Cx HCs, Panx channels, or P2X<sub>7</sub> receptor channels [72,73,94,171,176–180]. The downstream effect of these messengers involves the GPCR-PLC-IP<sub>3</sub> signaling axis or  $\text{Ca}^{2+}$  entry via plasma membrane



**Fig. 2.** Mechanisms underlying intercellular  $\text{Ca}^{2+}$  signaling. The activation of plasma membrane GPCRs results in the PLC-mediated synthesis of  $\text{IP}_3$  and subsequent IICR from the ER. Mitochondrial  $\text{Ca}^{2+}$  uptake prevents  $\text{Ca}^{2+}$  from reaching deleterious levels in the cytoplasm. The intracellular  $\text{Ca}^{2+}$  signal can propagate to neighboring cells via two routes, a direct one involving GJs, with  $\text{IP}_3$  as the primary coordinating messenger, and an indirect one that is mediated by the release of paracrine messengers. Here, ATP and glutamate are the prototypical paracrine molecules released into the extracellular environment via multiple mechanisms including HCs, a vesicular pathway or channels such as the  $\text{P2X}_7$  receptor. They diffuse in the extracellular space and activate their corresponding GPCRs or channels on neighboring cells. Regenerative mechanisms include the  $\text{Ca}^{2+}$ -dependent activation of  $\text{PLC}\delta$  or stimulation of paracrine release mechanisms. (CxHC, connexin hemichannel; ER, endoplasmic reticulum; GJ, gap junction;  $\text{G}_q\text{PCR}$ , G-protein coupled receptor; IICR, inositol 1,4,5-trisphosphate-induced  $\text{Ca}^{2+}$  release;  $\text{PIP}_2$ , phosphatidylinositol 4,5-bisphosphate; PLC, phospholipase C;  $\text{IP}_3$ , inositol 1,4,5-trisphosphate;  $\text{IP}_3\text{R}$ , inositol 1,4,5-trisphosphate receptor).

channels. Cx43 HCs may also contribute to the release of  $\text{NAD}^+$  that is converted extracellularly into cADPR. The plasma membrane CD38 enzyme subsequently transports cADPR back into the cell [76]. In most cases, ICW propagation is sustained by both the GJ and paracrine pathway, supported by regenerative steps which result in more extensive ICWs [9]. GJ-mediated regenerative processes rely on  $\text{Ca}^{2+}$ -triggered regeneration of  $\text{IP}_3$  via the activation of  $\text{PLC}\delta$  [163]. Paracrine messenger regeneration is supported by the  $\text{Ca}^{2+}$ -dependence of vesicular but also HC release mechanisms [86,181–183].

#### 4.2. Intercellular $\text{Ca}^{2+}$ signaling in the NGVU

Although several tissues and organs display ICWs or in more general terms ‘intercellular  $\text{Ca}^{2+}$  signaling’ (as the typical concentric wave aspect is not always present), the case is especially strong for the brain, where  $\text{Ca}^{2+}$  is a basic signal for glial network communication (Fig. 3). Evidence demonstrating  $\text{Ca}^{2+}$  signals propagating over tens to hundreds of astrocytes in cell culture started to appear more than 20 years ago [169–171,179,184–187]. Although the true function of inter-astrocyte  $\text{Ca}^{2+}$  signals is still unclear, the central position of astrocytes in the NGVU, making contacts with neurons, vascular cells and microglia (Fig. 3) has been a major factor in driving a continued interest and experimental exploration. There is now evidence for glial ICWs in more complex *ex vivo* preparations such as hippocampal and cortical slices [50,174,188–191]. For example, Weissman and colleagues demonstrated ICWs in radial glia that postnatally transform into astrocytes in the proliferative cortical ventricular zone of the developing neocortex [188]. The latter require Cx HCs,  $\text{P2Y}_1$  receptors and IICR, and are proposed to regulate cortical neuron proliferation. More recently, advances made in the field of two-photon microscopy have enabled visualization of astrocyte  $\text{Ca}^{2+}$  dynamics (not necessarily waves) *in vivo* (Table 3 and [16,17,21,192–194]), either arising spontaneously [16,17] or in response to somatosensory [195] or visual stimulation [194]. Spontaneously occurring wave-like  $\text{Ca}^{2+}$  signal propagation has been observed in glial cells of the hippocampus [18], the neocortex [17], the cerebellum


[16,19] and the retina [196] (Table 3). Local application of bicuculline to increase neuronal activity by suppressing GABA-signaling triggers astrocytic ICW activity *in vivo* [193] but evidence supporting astrocytic ICW generation in response to physiological levels of stimulation is still lacking.

Astrocytic ICWs have been reported to spread at a uniform speed of 5–20  $\mu\text{m/s}$ , which is several orders of magnitude slower than the propagation of neuronal electrical signals [16,188]. *In vivo*, ICWs propagate over astrocyte networks with even lower velocities in the order of 8–10  $\mu\text{m/s}$  [197]. The relative contribution of paracrine/purinergic signaling and GJIC to ICWs is likely to depend on brain region-specific factors [198] and on the nature of the applied trigger [188]. In acutely isolated retina, mechanical stimulation evokes ICWs among astrocytes which depend on GJIC, while the propagation from astrocytes to Müller cells involves purinergic, paracrine signaling [199]. In the corpus callosum and hippocampus, glial ICWs were blocked by purinergic receptor antagonists, but were insensitive to GJ blockers and the deletion of Cx43 or Cx30 expression using conditional Cx43(f/f):GFAP-cre mice and Cx30<sup>−/−</sup> KO mice [198,200]. In contrast, in the neocortex, glial ICWs depend on functional GJs as well as ATP release [198]. There is cross-talk between the gap junctional and the paracrine pathways of ICW propagation: deletion of Cx43 in spinal cord astrocytes leads to an increase in paracrine, purinergic  $\text{Ca}^{2+}$  signaling by a switch in the  $\text{P2Y}$  receptor subtype, resulting in increased responses to ATP or uridine 5'-triphosphate [102]. Additionally, the cytoplasmic C-terminal domain of Cx43 modulates the activity of the  $\text{P2Y}_1$  receptor in astrocytes via a protein–protein interaction between Cx43 and the  $\text{P2Y}_1$  receptor, involving the Cx43 C-terminal SH3-binding domain [101].

Astrocytic  $[\text{Ca}^{2+}]_i$  changes modulate synaptic transmission in *in vitro* astrocyte–neuron co-culture systems, the intact retina and hippocampal slices [61,201–203]. Astrocytes remove and recycle  $\text{K}^+$  ions and neurotransmitters from the synaptic cleft [204], but also respond to neurotransmitters by means of a large family of membrane receptors [37,192,205]. The ensuing signaling, often involving  $[\text{Ca}^{2+}]_i$  changes, triggers the release of gliotransmitters such as glutamate, D-serine,

ATP, adenosine and GABA [31–34,178,206–209]. These influence pre-synaptic neurotransmitter release, excitatory postsynaptic firing and extrasynaptic neuronal activity [207,210,211], neuronal circuit plasticity [212], and neuronal survival (see Section 5). The source of cytosolic  $\text{Ca}^{2+}$  that mediates gliotransmitter release is dual; it is predominately

derived from the ER involving both  $\text{IP}_3$ Rs and RyRs [213], but also involves  $\text{Ca}^{2+}$  influx from the extracellular environment, likely mediated by TRPC1 channels [214]. Additionally, mitochondria modulate exocytotic glutamate release from astrocytes, since inhibiting mitochondrial  $\text{Ca}^{2+}$  accumulation increases  $[\text{Ca}^{2+}]_i$  and enhances glutamate release



**Fig. 3.**  $\text{Ca}^{2+}$  signaling at the NGVU. At the level of the synapse, astrocytes perform three key functions. By removing and recycling byproducts of action potential firing, including glutamate and  $\text{K}^+$  ions, from the synaptic cleft, astrocytes protect neurons from excitotoxicity. Additionally, astrocytes are activated by neurotransmitters like glutamate to which they respond with an increase in  $[\text{Ca}^{2+}]_i$ . Subsequently, astrocytes release so-called gliotransmitters via which they influence synaptic activity and neuronal circuit plasticity. The increase in  $[\text{Ca}^{2+}]_i$  can be propagated to neighboring astrocytes via either the gap junctional or the paracrine route which will allow astrocytes to modulate remotely located synapses. The  $\text{Ca}^{2+}$  signal raised by synaptic activity will additionally propagate within the astrocytes toward the astrocytic endfeet that surround the blood vessels. Here,  $\text{Ca}^{2+}$  will stimulate the release of vasodilators and/or -constrictors. As such, astrocytes are important players in neuron-to-vessel signaling. Again, the propagation of  $\text{Ca}^{2+}$  signals between well-coupled endfeet will allow a coordinated response along the vessel wall. Finally, in pathological conditions, astrocytic  $\text{Ca}^{2+}$  signals may be involved in the activation of microglia. ATP release, possibly via HCs, is of significant importance in this process. Microglia may also be activated by damage-associated molecules, all of which lead to a microglial  $\text{Ca}^{2+}$  increase. This rise in  $[\text{Ca}^{2+}]_i$  may ultimately lead to the release of pro-inflammatory molecules that will signal back to astrocytes, contributing to astrogliosis. (20-HETE, 20-hydroxyeicosatetraenoic acid; GJ, gap junction; GLAST, glutamate/aspartate transporter; GLT-1, glutamate transporter 1; HC, hemichannel;  $\text{IP}_3$ , inositol 1,4,5-trisphosphate; mGluR, metabotropic glutamate receptor;  $\text{K}^+$ , inwardly rectifying  $\text{K}^+$  channel; NKCC,  $\text{Na}^+/\text{K}^+/\text{Cl}^-$  cotransporter;  $\text{Na}^+/\text{K}^+/\text{ATPase}$ ,  $\text{Na}^+/\text{K}^+/\text{ATPase}$ ; PGE<sub>2</sub>, prostaglandin E<sub>2</sub>; PRR, pattern recognition receptor; ROS, reactive oxygen species).

[215]. Notably, the  $\text{Ca}^{2+}$ -dependent glutamate release from astrocytes and its subsequent modulation of synaptic activity is challenged by a study performed on  $\text{IP}_3\text{R2}$  KO mice [216]. Several lines of evidence suggest that this subtype is the primary  $\text{IP}_3\text{R}$  expressed by astrocytes [217, 218]. Ablating  $\text{IP}_3\text{R2}$  eliminates both spontaneous and GPCR-mediated increases in astrocytic  $\text{Ca}^{2+}$ , while neuronal GPCR-dependent  $\text{Ca}^{2+}$  increases remain intact. However, silencing astrocyte  $\text{Ca}^{2+}$  signaling provoked by  $\text{IP}_3\text{R2}$  KO did not significantly affect perisynaptic glutamate levels nor did it alter excitatory CA1 pyramidal neuronal synaptic activity [216].

At the vascular side, astrocytic  $\text{Ca}^{2+}$  signaling links to blood flow regulation, metabolic trafficking, BBB function and water homeostasis [50]. In situ, astrocytes express  $\text{P2Y}_2$  and  $\text{P2Y}_4$  receptors and Cx proteins, both involved in  $\text{Ca}^{2+}$  signaling, as well as aquaporin 4 (AQP4) involved in water homeostasis at their perivascular endings [50]. This extraordinary high expression of basic  $\text{Ca}^{2+}$  signaling elements supports wave-like propagation of  $\text{Ca}^{2+}$  signals between rows of endfeet to influence vascular SMCs and blood vessel diameter, as observed in brain slices [219]. At present, it is not known whether endfeet  $\text{Ca}^{2+}$  signaling also influences capillary endothelial cells and BBB function. Cell culture studies have suggested propagation of ICWs between astrocytes and endothelial cells [68,220] but this could not be confirmed in cortical slice studies [50]. Possibly, this is related to the presence of the basal lamina interposed between endothelial cells and astrocytes, and the fact that ATP is a short-distance messenger that is rapidly degraded by ectonucleotidases [221].

The evidence that neuronal activity-triggered astrocytic  $\text{Ca}^{2+}$  signaling modulates arteriolar blood flow is particularly strong and supported by brain slice as well as in vivo animal studies [194,222–224], although at least one recent study challenges this view [225]. Electrical stimulation of neuronal afferents results in the activation of astrocytic metabotropic glutamate receptors (GluRs), giving rise to a  $[\text{Ca}^{2+}]_i$  increase and release of  $\text{PGE}_2$  that induces a vasodilation of adjacent cortical arterioles [223,224,226]. Follow-up work indicated that the  $[\text{Ca}^{2+}]_i$  increase in astrocytes is pursued by  $[\text{Ca}^{2+}]_i$  changes in vascular SMCs: vasodilation was accompanied by a suppression of  $\text{Ca}^{2+}$  oscillations in SMCs, likely mediated by astrocyte-derived carbon monoxide. The SMC response is furthermore abolished in isolated arterioles where astrocytes are absent and in brain slices where astrocytes are ablated by the gliotoxin L- $\alpha$ -aminoadipic acid (L-AAA), that is reported to target astrocytes by a yet unknown mechanism [38,227–229]. Astrocytic  $[\text{Ca}^{2+}]_i$  changes have, in addition to triggering vasodilation, also been reported to provoke vasoconstriction mediated by the arachidonic acid metabolite 20-hydroxyeicosatetraenoic acid [219]. One explanation for these opposite vascular reactions relates to perivascular  $\text{K}^+$  levels. The threshold concentration of perivascular  $\text{K}^+$  for inducing a vasodilatory response is 20 mM. When  $[\text{K}^+]$  exceeds this threshold, the vascular response converts into a constrictive one. Large conductance  $\text{Ca}^{2+}$ -sensitive  $\text{K}^+$  channels, that are strongly expressed in astrocytic endfeet [230], are decisive in determining these perivascular  $\text{K}^+$  levels [231,232]. Another explanation is related to oxygen availability. The vasodilatory response to astrocytic  $[\text{Ca}^{2+}]_i$  elevation is favored by low oxygen availability, which results in increased astrocytic glycolysis and consequent lactate release. This stimulates the extracellular accumulation of vasodilatory  $\text{PGE}_2$  and adenosine [233]. Astrocytic lactate release may also result from the uptake of synaptically derived glutamate, thereby boosting glycolysis and contributing as a  $\text{Ca}^{2+}$ -independent astrocytic influence [226]. A differential vascular effect of astrocytic  $[\text{Ca}^{2+}]_i$  elevation may be responsible for an increased blood flow in areas experiencing neuronal activation while reducing the flow to inactive regions vascularized by vessels branching from the same artery. Importantly, the extent of vasomotor responses seems to depend on the number of endfeet that are activated [222]. Spatially confined  $[\text{Ca}^{2+}]_i$  changes confer a dimensionally restricted control over the arteriolar diameter [194,219,224, 234]. By contrast, a sufficiently large increase in  $[\text{Ca}^{2+}]_i$  is capable of inducing an ICW that propagates along foot processes enwrapping the vascular wall, thereby mediating a more expanded vascular response

[50,219] which might be involved in the vasodilation of upstream pial arteries [235]. The role of glial Cx channels in this process remains to be fully established but at least one report highlights the necessity of Cx43 in vasodilatory responses [59]. It is unknown whether this concerns endothelial or astrocytic Cx43 but vascular Cx37 and Cx40 at least did not appear to be involved.

Alterations in local blood flow in response to neuronal activity are known as functional hyperemia or neurovascular coupling, and secure an adequate supply of glucose and oxygen under conditions of increased demand. Neurovascular coupling is adjoined by coupling of neuronal activity to changes in energy metabolism that is compartmentalized in astrocytes and neurons. According to the classical view, glucose is the main fuel for both neurons and astrocytes. However, this view has been challenged by the astrocyte–neuron lactate hypothesis which states that the uptake of glucose predominantly takes place in astrocytes. Subsequently, lactate is produced by astrocytes in an activity-dependent, glutamate-mediated manner, and is transferred to active neurons, where it's been converted into pyruvate that fuels the Krebs cycle [236–239]. Additionally, coupling to transport over BBB endothelial cells, dubbed neurobarrier coupling, has also been proposed [238, 240]. Each of these processes may be regulated by changes in  $[\text{Ca}^{2+}]_i$  as well [241–245].

## 5. Aberrant $\text{Ca}^{2+}$ signaling in neurogliopathologies

Spontaneous ICWs have been observed in vitro in cultured astrocytes starting from the 90s [184]. However, it is speculated that such spontaneous ICWs are probably an artifact of cultured cells [246]. In particular, the intensity/duration of the stimuli required to evoke these ICWs in vitro is often in a supraphysiological range [9]. Although frequently observed in cultured cells, several authors did not observe ICWs under basal/physiological levels of neuronal activity in vivo [21, 22,25,194,195]. In contrast, a handful of publications have reported the occurrence of ICWs in vivo in several disease states such as ischemia [20], epilepsy [25,26], AD [21,22], brain trauma [23,24] and cortical spreading depression [247] (Table 3), suggesting that pathological conditions are more favorable toward the triggering and/or the propagation of ICWs. However, many important questions still remain: how are these glial ICWs initiated, how are they propagated (paracrine versus GJIC) and how do glial ICWs contribute to the pathogenesis and disease progression?

### 5.1. Pathological triggers as potent inducers of intercellular $\text{Ca}^{2+}$ waves

A decrease in  $[\text{Ca}^{2+}]_e$ , known to evoke ICWs [187,248], has been reported to occur in hypoglycemia, ischemia, epilepsy, and CNS trauma [249–252]. In addition, mild declines in brain interstitial  $\text{Ca}^{2+}$  levels may take place in response to neuronal stimulation [253,254]. Various trigger mechanisms for the low  $[\text{Ca}^{2+}]_e$ -induced ICWs have been proposed with the most plausible one pointing to CxHC mediated ATP release that acts as a point source from trigger cells [169]. Mechanical stimulation and trauma induce ICWs that can propagate hundreds of  $\mu\text{m}$  in in vitro model systems, depending on the magnitude of the stimulus [23,24,255,256]. Wave-like  $\text{Ca}^{2+}$  dynamics in vivo generally appear to involve tens of astrocytes [16,26,257]. Amyloid- $\beta$  (A $\beta$ )-peptide, the major component of senile plaques in Alzheimer patients, increases the amplitude, velocity and distance of ATP- and GJ-mediated ICWs in astrocyte cultures [258]. Moreover, pathologically relevant concentrations of A $\beta$  (nM to  $\mu\text{M}$  range) were able to induce astrocytic ICWs with a delay of minutes, suggesting a reorganization in terms of chemical  $\text{Ca}^{2+}$  excitability threshold and/or communication properties of the intercellular network [259]. Of note, picomolar amounts of A $\beta$  that are not associated with pathology, do facilitate spontaneous  $\text{Ca}^{2+}$  transients in astrocytes but do not affect ICW parameters [260], pointing to an increase in  $\text{Ca}^{2+}$  excitability. In amyloid precursor protein (APP)/presenilin 1 (PS1) double transgenic mice that express mutant human PS1- $\Delta\text{E9}$  and chimeric mouse/human APP<sub>swe</sub>, A $\beta$  plaques nearly double



**Table 3**Overview of triggers and mechanisms involved in in vivo intercellular  $\text{Ca}^{2+}$  signaling.

Trigger	Cell types <sup>a</sup>	Propagation mechanism	Remarks	Reference
<i>Spontaneous</i>				
	Astrocytes/neurons	ND	ICWs observed in astrocytic network	[17]
	Hippocampal astrocytes	Purinergic signaling/Cx channels	High-speed ICW-like propagations ('glissandi') involving hundreds of astrocytes	[18]
			Glissandi depend on neuronal activity	
			Associated with a reduction in blood flow	
	Retinal glia	Purinergic signaling	ICWs propagated through Müller cells, not through astrocytes	[196]
			Aging increases spontaneous ICWs due to more ATP release	
			Associated with a reduction in blood flow	
	Bergmann glia	ND	Aging increases spontaneous ICWs	[19]
			Accompanied by $[\text{Ca}^{2+}]_i$ increases in Purkinje neurons, but no change in neuronal activity	
			Reduces brain oxygen tension	
	Bergmann glia	Purinergic signaling	Recurring waves extend from the same foci	[16]
<i>Neuronal activity</i>				
Bicuculline (GABA <sub>A</sub> receptor blocker)	Astrocytes			[193]
Whisker stimulation	Astrocytes	Glutamatergic signaling	No ICWs reported	[195]
Visual stimuli	Astrocytes	Glutamatergic signaling	No ICWs reported	[194]
Glutamate uncaging/visual stimuli	Microglia	Neuronal Panx1 HCs/purinergic signaling	No ICWs reported	[384]
			Microglia migrate to and make contact with activated neurons	
			Microglial–neuronal contact downregulates neuronal activity	
<i>Ischemia</i>				
Photothrombosis	Astrocytes	Glutamatergic	The $\text{Ca}^{2+}$ chelator BAPTA-AM reduced the infarct volume	[20]
<i>Alzheimer's disease</i>				
APP/PS1 mice	Astrocytes	ND	Astrocytic $[\text{Ca}^{2+}]_i$ elevated, independently from distance to A $\beta$ plaques	[21]
			ICWs induced near plaques and spread radially	
3xTgAD mice	Astrocytes	ND	In contrast to AD mice, no spontaneous ICWs or ICWs in response to photo-activation of $\text{IP}_3$ were observed in control mice	[22]
PDAPP mice				
<i>Traumatic injury</i>				
Brain cortical impact	Astrocytes	Purinergic signaling	Influences neuronal network activity and evokes neuronal death as evidenced from experiments conducted on neuron–glia co-cultures and organotypic slice cultures	[24]
			ICWs not dependent on ATP or Cx channels	
			ICW-evoked ATP release attracts microglia	
			Local photo-activation of caged $\text{IP}_3$ mimicks the microglial injury response	
Laser ablation	ND	Glutamatergic signaling		[23]
			ICW-evoked ATP release attracts microglia	
			Local photo-activation of caged $\text{IP}_3$ mimicks the microglial injury response	
Laser ablation/mechanical injury	Astrocytes	Purinergic signaling/Cx channels	Polarized $\text{Ca}^{2+}$ gradient in astrocytes that guides microglia to the injury site	[376]
			No ICWs observed	
<i>Epilepsy</i>				
Pilocarpine-induced status epilepticus	Astrocytes	Glutamatergic signaling	Evoke neuronal cell death	[25]
			Astrocytic $\text{Ca}^{2+}$ signals are dependent on metabotropic mGluR5, while neuronal cell death relies on NMDAR activity	
			Evokes PDSs through glutamate release	
4-Aminopyridine-induced seizures	Astrocytes	ND		[26]
<i>Cortical spreading depression</i>				
High $[\text{K}^+]_e$	Neurons/astrocytes		ICWs are observed in neurons and in astrocytes	[247]
			Astrocyte ICWs evoke vasoconstriction of intracortical vessels	

<sup>a</sup> In which the  $\text{Ca}^{2+}$  responses were studied. (A $\beta$ , amyloid- $\beta$ ; APP, amyloid precursor protein; BAPTA-AM, 1,2-bis(2-aminophenyl)ethane-N,N,N',N'-tetraacetic acid acetoxymethyl ester; Cx, connexin; HC, hemichannel; ICW, intercellular  $\text{Ca}^{2+}$  wave; mGluR, metabotropic glutamate receptor; NMDAR, N-methyl-D-aspartate receptor; PDS, paroxysmal depolarization shift; PS, presenilins)

astrocyte resting  $[\text{Ca}^{2+}]_i$  and promote spontaneous  $\text{Ca}^{2+}$  activity that is communicated radially over at least 200  $\mu\text{m}$  along the astrocytic network [21]. The ICWs do not rely on the presence of hyperactive neurons near the plaques [21,261], indicating that the increased astrocytic  $\text{Ca}^{2+}$  excitability is induced by A $\beta$  exposure itself and is not the result of an enhanced electrical excitability of neurons. The augmentation in astrocyte  $[\text{Ca}^{2+}]_i$  in response to A $\beta$  can be mediated by  $\text{Ca}^{2+}$  store release ( $\text{IP}_3$ - and RyR-dependent) and SOCE [262]. Additionally, A $\beta$  has been shown to bind the  $\text{Ca}^{2+}$ -sensing receptor, a GPCR that senses extracellular  $\text{Ca}^{2+}$  levels [263], or to enhance spontaneous  $\text{Ca}^{2+}$  signaling via nicotinic acetylcholine receptors in astrocytes [260]. Alternatively, intracellular accumulation of  $\text{Ca}^{2+}$  may be mediated by the pore-forming capacity of A $\beta$  protein or possibly by creating a non-selective  $\text{Ca}^{2+}$  entry route by inducing oxidative stress and lipid peroxidation of the plasma membrane [264,265].

Another potential trigger for ICWs may originate from areas of cell death. Most, if not all, brain diseases are associated with cell death

that is accompanied by the release of ATP and glutamate from the intracellular metabolic pool [266]. These neural/glial transmitters activate ionotropic and metabotropic receptors, with subsequent  $\text{Ca}^{2+}$  entry and  $\text{IP}_3$ -triggered ER  $\text{Ca}^{2+}$  release that may trigger ICWs [169–173, 184,186]. Work from our research group on cultured glioma cells, has brought forward that both Cx43 GJs and HCs are involved in communicating cell death messages from cells undergoing apoptosis to healthy neighboring cells, a process known as 'bystander cell death' [267]. Importantly, we demonstrated that the passage of the  $\text{Ca}^{2+}$  messenger  $\text{IP}_3$  through GJs is crucial for intercellular cell death communication. This work provided evidence that the physiological messenger  $\text{IP}_3$  becomes a pathological intercellular messenger when the cells generating  $\text{IP}_3$  are exposed to proapoptotic conditions [268,269]. Recent work from others supports this concept and demonstrates that HIV infection of astrocytes results in bystander apoptosis of uninfected astrocytes via the  $\text{IP}_3/\text{Ca}^{2+}$  signaling axis [270]. Thus, IICR is required for normal function and cell survival, but it can also be actively involved in apoptosis

induction, progression and propagation [269,271]. In fact, ER-mitochondrial  $\text{Ca}^{2+}$  transfer at the level of the MAMs plays a major role in cell death by providing proapoptotic  $\text{Ca}^{2+}$  transfer via  $\text{IP}_3\text{R}$  interaction with VDAC1 [143,272–276]. Additionally, numerous factors involved in apoptosis have been located at the ER-mitochondrial interface and modulate cell survival by directly or indirectly regulating IICR [277,278]. Recent evidence revealed that alterations in ER-mitochondrial  $\text{Ca}^{2+}$  communication, possibly by regulating IICR, may represent a common hit in several brain pathologies including ischemia [279,280] and neurodegenerative diseases such as AD and Parkinson's disease [150,281–286]. Although neurons themselves are the primary target of investigation in these studies, an increasing number of publications have demonstrated alterations in ER  $\text{Ca}^{2+}$  signaling in astrocytes, primarily in ischemic conditions and AD [280,287–290]. While the contribution of GJs to bystander cell death is straight forward, i.e. they facilitate the direct exchange of cell death and cell survival signals between cells, the role and functional consequences of HC opening on cell death is still poorly defined. Several possibilities have been put forward: (i) HC opening may help in setting off cell death by the excessive entry of  $\text{Ca}^{2+}$  or other toxic molecules and (ii) being a pore permitting diffusion in both directions, HCs could also contribute via the release of survival molecules or, more indirectly, via the release of paracrine messengers which induce a  $[\text{Ca}^{2+}]_i$  rise in more distant cells [291,292].

Disturbances in astrocyte  $\text{Ca}^{2+}$  signaling combined with Cx-mediated  $\text{Ca}^{2+}$  signal communication may thus act to expand cell death and neurodegeneration. As discussed in the next sections, ICWs may possibly be involved in expanding the processes of reactive astrogliosis and microglial activation as well. The balance between astrogliosis, microglial activation and neurodegeneration may ultimately determine the level of destruction, neuroprotection and regeneration resulting from the pathological process.

### 5.2. A role for $\text{Ca}^{2+}$ and connexins in astrogliosis?

Reactive astrogliosis, a defensive astrocytic response, is a common feature of many CNS disorders, including stroke, trauma, epilepsy, neurodegeneration, infection and autoimmunity [293]. However, despite its widespread observation, the role of as well as the cellular changes accompanying astrogliosis are poorly understood. Reactive astrogliosis is generally regarded as a defensive reaction towards injury which aims at isolating the damaged area from the rest of the brain tissue to facilitate restoration of the tissue surrounding the lesion. It generally appears as a hypertrophy and/or proliferation of astrocytes surrounding the lesion. Based on the severity of the changes, reactive astrocytes can be subdivided in two different classes, anisomorphic and isomorphic [294]. Anisomorphic astrocytes are characterized by robust hypertrophy and proliferation and evolve towards the formation of glial scar tissue that is composed of astrocytes and extracellular matrix [295]. Glial scar tissue surrounds the injured/lesioned tissue site and may therefore prevent its extension but it also plays a key role in regeneration failure because it inhibits axonal sprouting. In isomorphic astrocytes, the reactive changes are milder and often reversible, allowing regeneration of the tissue.

The upregulation of intermediate filament proteins, including glial fibrillary acidic protein (GFAP) and to a lesser extent nestin and vimentin, are often used as indicators for astrogliosis [296]. Other changes may appear in a context/disease-specific manner [297]. It is important to realize that many proteins associated with  $\text{Ca}^{2+}$  signaling are altered in reactive astrocytes. Genetic profiling of the astrocyte response towards pro-inflammatory mediators such as TGF- $\beta$  and IFN- $\gamma$  has recently uncovered the specific responses of different GPCRs towards inflammation and has revealed pronounced changes in the  $\text{Ca}^{2+}$  household during astrogliosis [298]. Likewise, an upregulation of the  $\text{Ca}^{2+}$ -entry channel TRPV4 coincides with the onset of astrogliosis during an hypoxic/ischemic insult [299], whereas the L-type  $\text{Ca}^{2+}$  channel  $\text{Ca}_v1.2$  is upregulated in reactive astrocytes surrounding A $\beta$  plaques

[300]. Most interestingly, in several models,  $\text{Ca}^{2+}$  appears to be one of the early injury signals that initiates the signal transduction cascade for astrogliosis; e.g. through ER  $\text{Ca}^{2+}$  store release (via  $\text{IP}_3\text{Rs}$  and RyRs) and SOCE in A $\beta$ -treated astrocytes [259,262] or following lipopolysaccharide (LPS) exposure [301], and through IICR [302] and  $\text{Ca}^{2+}$  influx in response to spinal cord and brain injury [303,304].

Cx43 is also reported to be one of the proteins modified in reactive astrocytes. Increased Cx43 expression has been detected in regions with astrogliosis induced by various CNS diseases including brain injury [305–308], spinal cord injury [309,310], brain ischemia [311,312], AD [313], epilepsy [314,315], and also following retinal ischemia [316, 317]. Cx43 deletion in astrocytes causes a reduction of astrogliosis in brain ischemia and brain/spinal cord injury models [306,309, 318–321]. Inversely, astrocytic Cx43 was decreased in the spinal cord of experimental autoimmune encephalomyelitis (EAE) mice (a model for multiple sclerosis [322]), and in response to a cortical lesion induced by kainic acid injection [323]. Thus, Cx43 may be up- or down-regulated, depending on the type of brain pathology, but probably also on the time scale, and the distance from the lesioned area. For example, excitotoxic induction of neuronal cell death *in vivo* lead to a strong decrease of Cx30 and Cx43 in reactive astrocytes located within the neuron-depleted zone, whereas increased Cx expression was found in reactive astrocytes at the periphery of the lesion [51]. Transient Panx2 expression was observed in reactive astrocytes in the hippocampus of adult rat brain subjected to ischemia/reperfusion injury [324].

There is limited evidence that ICWs could contribute to the generation of reactive astrocytes and glial scars e.g. after traumatic injury [255], ischemia [20], and A $\beta$  exposure [259]. One paper has demonstrated an involvement of Cx-mediated ICWs to the expansion of astrogliosis [304]. Scratch injury applied to mouse cortical astrocyte cultures induced a  $[\text{Ca}^{2+}]_i$  elevation along the scratch via  $\text{Ca}^{2+}$  entry, which was then disseminated to neighboring astrocytes as an ICW propagating via GJs. These increases in  $[\text{Ca}^{2+}]_i$  switched on the JNK/c-Jun/AP-1 pathway to upregulate GFAP expression. Dampening the  $[\text{Ca}^{2+}]_i$  changes with intracellularly-loaded BAPTA in an *in vivo* stab wound model, reduced the expression of GFAP and the extent and size of glial scars, supporting the involvement of  $\text{Ca}^{2+}$  signaling [304].

### 5.3. Neuronal cell death, a consequence of astrocyte network distortion?

Excitotoxicity, induced by excessive glutamate release and subsequent activation of  $\text{Ca}^{2+}$  and ROS signaling, is a major trigger for neuronal cell death [325,326]. Astrocytes coupled by GJs normally act to promote neuronal survival after a pathological insult by scavenging ROS (astrocytes contain high concentrations of antioxidant molecules such as glutathione and ascorbate) and by spatial buffering of extracellular  $\text{K}^+$  and glutamate as set out in Section 3. However, as things get rough, glial homeostatic mechanisms can be altered which rapidly provides the astrocytic network with killer functions including the exacerbation of neuronal damage by decreased glutamate or  $\text{K}^+$  uptake or by increased release of glutamate, ATP or inflammatory cytokines [319, 327–331]. Most notably, ICWs spreading over the astrocytic network may promote astrocytic release of these messengers thereby worsening neuronal damage. The dual role of astrocytes implicates that a precise regulation of their response may determine the outcome of CNS pathology.

Astrocytic ATP release combined with extracellular glutamate accumulation promotes neuronal cell death [327] and at millimolar extracellular ATP concentrations it contributes to astrocytic cell death via the activation of P2X<sub>7</sub> receptors [332]. On the other hand, ATP is rapidly degraded to adenosine, which can affect neuronal functioning as it inhibits excitatory activity and presynaptic neurotransmitter release [254,333]. ATP may also contribute to myelin disturbances and axonal degeneration; ischemia-induced oligodendroglial death shows hallmarks of a  $\text{Ca}^{2+}$ -dependent excitotoxic pathway which involves the activation of N-methyl-D-aspartate receptors (NMDARs) and P2X<sub>7</sub>

receptors that are activated by ATP released through Panx channels [334]. Panx1 channels were also suggested to function as large pores associated with the P2X<sub>7</sub> receptor [335,336], including in astrocytes [93]. Panx1 co-immunoprecipitates with P2X<sub>7</sub>R and inhibition of Panx1 prevents P2X<sub>7</sub>R-mediated dye uptake [335,336]. In addition, functional NMDAR have been identified in astroglial cells (reviewed in [337]) and excess activation results in a Ca<sup>2+</sup> influx that may spread through GJs composed of Cx43 and Cx47 to oligodendroglia, evoking apoptosis in the latter [338]. Finally, astrocytes produce and release a wide variety of inflammatory molecules including cytokines, chemokines and prostaglandins [339–341], which provides a mutual communication pathway between astrocytes and microglia (see Section 5.4). Astrocytes may be involved in anti-inflammatory processes including scar formation [342], or pro-inflammatory cytokine release may damage cells by e.g. increasing the Ca<sup>2+</sup>-dependent release of glutamate [343]. Among the cytokines expressed by astrocytes, interleukin-1 $\beta$  (IL-1 $\beta$ ) and tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) have received considerable attention because they activate CxHCs and exert pro-inflammatory effects in the CNS [331,344]. Inhibition of Panx1 prevents activation of innate immunity and secretion of pro-inflammatory cytokines through the pattern recognition receptors NLRP1 and NLRP2 [345,346]. Panx1 channels, P2X<sub>7</sub> receptors and CxHCs may thus act in concert during the course of neuroinflammation.

#### 5.4. Neuroinflammation, a combined effort between activated astrocytes and microglia

Ischemia, trauma and neurodegenerative diseases such as AD, amyotrophic lateral sclerosis (ALS), Huntington's disease and Parkinson's disease are some of the pathologies which are accompanied by inflammatory events [347,348]. Microglial cells form the innate immune defense system of the CNS. They continuously scan the brain parenchyma for damage signals by which they are activated. Once activated, microglial cells migrate to the site of injury, proliferate, and phagocytize cells and cellular compartments [349]. They accumulate around degenerating neurons and reactive astrocytes in various CNS disorders [347,348] and show diverse activated phenotypes depending on the pathology and the damage signal or chemoattractant that is released [350,351]. The best known chemoattractant sensed by microglia is ATP [352]. Several neuropeptides such as angiotensin II, bradykinin, endothelin, galanin and neurotensin are also chemoattractants for microglia [353]. Most of these molecules activate microglia by triggering a [Ca<sup>2+</sup>]<sub>i</sub> elevation via plasma membrane Ca<sup>2+</sup> channels such as ionotropic P2X receptors, IICR and the reverse mode of the NCX [351,354]. Microglia can also be activated by an A $\beta$ -induced [Ca<sup>2+</sup>]<sub>i</sub> surge, after which they secrete pro-inflammatory and neurotoxic factors [355,356].

Activated microglia release a large number of substances including ATP, glutamate, cytokines like TNF- $\alpha$ , superoxide radicals and neurotrophins that can either act detrimental or beneficial for the surrounding cells. They communicate with neurons and macroglial cells through paracrine factors [357]. Activated microglia express Cx32, Cx36 and Cx43 [357]. It is conceivable that the paracrine signaling molecules, some of which have a MW < 1.5 kDa, may be released via HCs. There is in vitro evidence for microglial glutamate release via HCs upon exposure to TNF- $\alpha$  [358]. Blocking HC-mediated glutamate release via different pharmacological substances (see Section 5.5) furthermore protects neurons from ischemic brain injury, attenuates EAE, and suppresses disease progression in ALS and AD mice in vivo [359–361]. Astrocytes can rescue neurons from microglial glutamate-induced death [362] but their glutamate-uptake capacity can also be hampered by microglial glutamate release via downregulation of the astrocytic GLAST glutamate transporter, thereby worsening extracellular accumulation of glutamate [363]. Cx32 HCs have been implicated in the release of glutamate from microglia [358]; however, at present, evidence for the expression and functional role of Cx32 remains poor. HCs composed of Cx43 are other candidates for mediating the release of paracrine

messengers. This Cx is not expressed in quiescent microglial cells, but it is upregulated upon their activation, suggesting a role during inflammation. It has been recently shown that ATP release via microglial Cx43 HCs is involved in microglial death [364]. Finally, although GJs have been described in cultured microglial cells [365], GJIC between microglial cells as well as with other surrounding cells has recently been reported to be absent in vivo [366].

Conditioned media from activated microglia or direct application of a mixture of the pro-inflammatory cytokines TNF- $\alpha$  and IL-1 $\beta$  influence the Cx-based astrocytic communication network: GJIC is reduced while HCs are invariably reported to open [78,331,344,367–369]. Effectwise, the reduced GJ coupling will restrict the movement of glucose and lactate over the astrocytic network while the open HC pathway may provide non-specific glucose uptake from the interstitium [49,78]. The net balance of these diverse effects is currently not clear, but they might have serious implications for the metabolic status of astrocytes in brain inflammation. In terms of ICWs, pro-inflammatory cytokines can shift the ICW supportive pathway from GJIC to paracrine mechanisms [370]. Cytokines released by astrocytes can also influence microglial Ca<sup>2+</sup> signaling [371] and were demonstrated to increase GJIC and induce HC opening in microglial cells [372,373].

[Ca<sup>2+</sup>]<sub>i</sub> increases evoked by danger signals can propagate as ICWs, thereby possibly expanding inflammation. A recent study in zebrafish larval brain revealed that laser light-induced neuronal injury initiates an ICW that travels throughout the brain and attracts microglia to migrate to the injury site. Here, glutamate-dependent ICWs establish an ATP guidance cue that is responsible for transmitting positional information across the brain to P2Y<sub>12</sub>-expressing microglia [23]. The propagation of the ICWs did not depend on Cx channels and ATP signaling, which contrasts with the observation that chemotactic microglial activation in mice is reduced by Cx channel blockers [374]. Although neurons, oligodendrocytes and endothelial cells are all candidates for releasing ATP upon injury, astrocytes are suggested to be the main cell type responsible for the rapid and long-range microglial response towards injury. Astrocytic ICWs trigger microglial responses in brain slices which appear to depend on the release of ATP [200,375]. Kim and Dustin [376] reported that laser-induced necrotic injury in vivo induces morphological changes in astrocytes that were accompanied by a [Ca<sup>2+</sup>]<sub>i</sub> gradient (but not ICWs) in the direction of the injury site. The astrocyte polarization and Ca<sup>2+</sup> response was proposed to guide microglial cells and was blocked by ATP and Cx channel blockers.

Panx1 channels have also been extensively studied as ATP releasing pores. They seem to play a key role in the release of this “find-me” signal from apoptotic cells for phagocytotic clearance [377,378]. Panx1 channels can be activated in cultured neurons and astrocytes, and in hippocampal slices, by elevated [K<sup>+</sup>]<sub>e</sub>, which can occur during different pathological conditions including epilepsy [379,380]. They have been proposed to mediate astrocytic ICWs via the release of ATP [94,181]. In leeches, nerve injury induces ICWs in glial cells and ATP release through channels composed of innexins, the invertebrate analogues of Cxs. Extracellular ATP then activates microglia and initiates microglial movement [256]. The injury site is also characterized by a delayed release of arachidonic acid, which blocks ATP release from glia after nerve injury and thereby stops microglia at lesions [381]. Panx1 has furthermore been implicated in the activation of the inflammasome, a multiprotein complex involved in innate immunity, in astrocytes and neurons [379]. Moreover, a number of studies have supported the idea that Panx channels are the pore-forming component of the P2X<sub>7</sub> receptor, with prolonged stimulation resulting in cell death [334,335,382]. Of note, from several studies it is also clear that extracellular ATP release not only mediates a rapid microglial response towards injury but it also regulates microglial branch dynamics in the intact healthy brain [374,383]. By imaging the visual center optic tectum in living larval zebrafish, Li and co-workers recently demonstrated a bi-directional regulation between resting microglia and neurons: a local increase in



neuronal activity results in the neuronal release of ATP via Panx1 channels, which attracts resting microglial processes and drives them to contact neurons with high levels of activity. The microglia–neuron contact in turn reduces the activity of contacted neurons [384].

### 5.5. Connexin-mediated communication, gap junctions versus hemichannels

Cx-mediated intercellular communication has been suggested to support the propagation of both pro-survival and pro-cell death messengers [291]. While treatments that suppress Cx43-based communication can re-inforce neurotoxicity in some models of neurodegenerative conditions [385,386], they may also be neuroprotective [318,319,387,388]. These opposite functions could relate to differences in the model systems used [291]. It should be noted that many of these observations were based on the use of non-specific channel blockers and/or genetic silencing of Cx expression, which affect both GJ and HC functioning. In addition, the genetic silencing of Cxs can modulate the expression levels of other proteins. It was recently demonstrated that the Cx43 and Cx30 deficiency in double KO mice differentially affects the expression of glial glutamate transporters depending on the brain region studied [389]. As such, more appropriate experimental tools are crucial to gain insight into the mechanisms via which Cxs affect cellular injury, e.g. by applying specific blockers that affect HCs or GJs only. Certain peptides derived from the Cx43 protein sequence can act as selective inhibitors of Cx channels [390–392]. Peptides targeting the extracellular loops that inhibit HCs within minutes and GJs after several hours [393], are available for Cx26, Cx32, Cx37, Cx40 and Cx43 [392,394,395]. The Cx mimetic peptides Gap26 and Gap27, derived from the Cx43 sequence, have been used extensively to inhibit Cx channels in glial cells, giving rise to various effects such as inhibiting glutamate and ATP release, protecting neurons from excitotoxicity, inhibiting apoptosis and influencing neuroinflammation [86,267,331,344,396]. In addition, other peptides identical to sequences on the intracellular parts of the Cx43 protein have been exploited to specifically inhibit HCs without influencing GJs [390,397,398]. Cx43 targeting peptides have been demonstrated to resort to beneficial effects in in vitro and in vivo models of CNS pathologies, including brain and retina ischemia [316,399,400], epilepsy [388,401], AD [402], optic neuropathy [403], spinal cord injury [319,404], and inflammation-induced BBB permeability [394].

### 5.6. Astroglial intercellular $\text{Ca}^{2+}$ waves, a common factor in neurodegeneration?

The vast majority of research tackling the question of the Cx/Panx channel contribution to cell death has been performed in the context of the CNS. An increasing number of reports demonstrate a profound effect of pathological conditions on the expression of glial Cxs and Panxs, in particular Cx43 and Panx1, and their function as channels in a range of CNS diseases including epilepsy, traumatic brain and spinal cord injury, AD, and ischemia-related injury [405–410]. Below, we discuss the role of ICWs and Cx/Panx alterations in each of these disease states.

#### 5.6.1. Brain ischemia

Cx-mediated intercellular communication may modulate the gradual expansion of cell death, as has been well-exemplified in the case of ischemia-induced injury. Although neurological cell death occurs within seconds to minutes after the ischemic insult in the core region, the progression of ischemic injury and cell death can continue for hours and days, thereby expanding the infarct region to the surrounding area termed the penumbra [411–414]. The latter is characterized by the presence of viable, yet metabolically compromised, cells that can be rescued by either restoring/improving perfusion or by counteracting neuronal damage. Many conceptually promising, therapeutic strategies based on targeting a variety of molecular cell injury/cell death pathways in neurons have failed in human studies, mainly because messengers causing destruction are also necessary for remodeling and repair

afterwards [415]. Thus, specific messengers leading to cell injury need to be targeted at the right time before they adopt beneficial roles. Additionally, focus has shifted from the neurons towards the astrocytes, which are considered today as novel therapeutic targets [416–418]. Astrocytes are more resilient to ischemic injury compared to neurons, most likely due to their high levels of antioxidants and their prominent anaerobic energy metabolism [237,419]. However, once the astrocytic network fails, glutamate/ $\text{K}^+$  clearance and energy delivery through the astrocyte–neuron lactate shuttle will stop and active release of neurotoxic mediators may start [235,236,239,420–422]. Hence, astrocyte-network distortion can critically contribute to neuronal death.

Cxs have been suggested to contribute to the wave of delayed secondary injury that spreads from the core region to the penumbra. Most studies emphasize a role for glial Cxs [408,423–426] but evidence for a contribution of neuronal GJs composed of Cx36 is accruing as well [42,427]. Ischemia-related injury is mostly associated with a reduction of astrocyte GJIC and an aberrant opening of Cx43 HCs [84,344,400,428,429]. However, despite the reduction in GJIC, there is evidence that a subset of GJs remains open and thus available for transferring cell death or survival signals under ischemic conditions [84,291,292,430]. Additionally, GJs may contribute to the propagation of cortical spreading depression (CSD), which consists of waves of tissue depolarization (20–80  $\mu\text{m/s}$ ) that have been linked to an increase in infarct volume after an ischemic insult [57,431–433]. CSD has been associated with slower propagating ICWs in astrocytes in vitro [190,434,435], and more recently also in vivo [247]. Here, CSD-associated ICWs first occurred in neurons but this was tightly coupled to a preceding ICW in astrocytes and to the vasoconstriction of intracortical arterioles [247]. The opening of Cx43 HCs will not only lead to astrocyte dysfunction and cell death by a collapse of membrane potential and the entry of  $\text{Ca}^{2+}$  [84,400,428,429], but will additionally allow the escape of glutamate and ATP that may further promote cell death [291,292,424].

Evidence for ICWs contributing to the expansion of brain infarction is also available. In response to photothrombosis-induced ischemia, cortical astrocytes in vivo showed large repetitive and transient  $\text{Ca}^{2+}$  signals within 20 min after photothrombosis which lasted up to 3 h [20]. They were synchronized and propagated as ICWs within the glial network, giving rise to  $\text{Ca}^{2+}$  signals in astrocytes in the penumbra. In contrast to neurons, astrocytes in the ischemic core retained their structural integrity for hours while an increase in reactive astrocytes was observed in the penumbra. Cortical loading of the  $\text{Ca}^{2+}$  buffer BAPTA into astrocytes significantly reduced the infarct volume. The astrocytic  $\text{Ca}^{2+}$  signals were triggered by mGluR5 and GABA<sub>B</sub> receptors [20]. A reduction of brain oxygen tension evokes increased frequency of spontaneous cerebellar ICWs in Bergmann glia in vivo, and  $[\text{Ca}^{2+}]_i$  increases in Purkinje neurons [19]. Several authors have reported a role for astrocytic IP<sub>3</sub>R<sub>s</sub> in modulating ischemic damage. IICR is shown to be neuroprotective since the ischemic damage was aggravated in IP<sub>3</sub>R2 KO mice which abolish astrocyte  $\text{Ca}^{2+}$  dynamics [218]. Finally, treatment with 2-aminoethoxydiphenyl borate, an inhibitor of IICR, decreased spontaneous  $\text{Ca}^{2+}$  oscillations in both astrocytes and neurons, and induced ischemic tolerance via a down-regulation of the glutamate transporter-1. A subsequent build-up of extracellular glutamate desensitizes the cells to the ischemia-induced glutamate increase [436]. An important issue to take into account is that 2-aminoethoxydiphenyl borate has off-target, inhibitory effects on Cx channels [437] and SOC<sub>s</sub> [438].

ICWs may be initiated by excessive neuronal firing or cell injury in the lesioned area. Once they are activated, they may self-sustain and expand  $[\text{Ca}^{2+}]_i$  increases over larger populations, resulting in an expansion of the damaged area. However, as evidenced by a recent paper, astrocyte  $\text{Ca}^{2+}$  signaling is not necessarily a bad thing. In response to an ischemic insult, astrocytes in the contralateral hemisphere experience a boost in  $\text{Ca}^{2+}$  signaling which contributes to enhanced uptake of glutamate, thereby eliminating the excitotoxic trigger [439]. As contralateral astrocytes are not subject to energy deprivation this glutamate may be converted to glutamine and shuttled to starving



neurons. As such, astrocyte  $\text{Ca}^{2+}$  signaling might have different functions/consequences depending on the proximity of the lesioned area. Similar concepts may apply to other CNS diseases characterized by damaged core regions such as AD (A $\beta$  plaques), brain trauma and epilepsy.

### 5.6.2. Alzheimer's disease

AD is considered as one of the most prevalent forms of dementia in humans. It is associated with several histopathological changes, including senile plaque formation, neurofibrillary tangles (NFTs), neuroinflammation, altered synaptic connectivity and ultimately irreversible, fatal neurodegeneration. NFTs are intraneuronal aggregates of paired helical filaments of abnormally hyperphosphorylated Tau protein. Senile plaques are extracellular aggregates mainly composed of A $\beta$  surrounded by dystrophic neurites, activated microglia, and reactive astrocytes. AD mostly occurs as a late-onset, idiopathic or non-Mendelian form, while only about 5% is familial. Familial AD has an early onset and is inherited in an autosomal dominant manner due to mutations in the amyloid pathway such as in APP or PS1/2, which are the catalytic subunit of the  $\gamma$ -secretase complex that cleaves APP and releases A $\beta$ .

The etiology of AD is poorly understood. The impairment of  $\text{Ca}^{2+}$  homeostasis is postulated to be an early event in the development of neurodegenerative processes [440]. Some suggest that altered  $\text{Ca}^{2+}$  signaling may decrease synaptic strength and cell loss well before initiating A $\beta$  plaques while others place A $\beta$  plaque formation as the start of  $\text{Ca}^{2+}$  deregulation [440,441]. The accumulation of NFTs and A $\beta$  plaques may alter the  $\text{Ca}^{2+}$  household of the nervous tissue. For example, Tau proteins alter the function of voltage-gated  $\text{Ca}^{2+}$  channels [442] whereas PS form  $\text{Ca}^{2+}$  leak channels in the ER [443]. PS physically interact with SERCA, IP $_3$ R and RyRs, facilitating ER  $\text{Ca}^{2+}$  uptake and release [283,444,445]. FAD-linked mutations in PS have been found to either abolish its  $\text{Ca}^{2+}$ -leak properties [443] or to result in a gain-of-function of IP $_3$ R channels [283]. The latter seems to activate a vicious circle, further increasing the ratio of the toxic A $\beta_{42}$  over A $\beta_{40}$  levels. Interestingly, A $\beta_{42}$  can directly perturb intracellular  $\text{Ca}^{2+}$  homeostasis via IP $_3$ R-dependent as well as -independent mechanisms [281]. A $\beta$  also compromises the ability of neurons to restore  $[\text{Ca}^{2+}]_i$  to baseline after a kainate and NMDA-induced elevation, making them more vulnerable to excitotoxicity [446]. In vivo evidence indicates that neurons in close proximity to A $\beta$  plaques are hyperactive and experience a  $\text{Ca}^{2+}$  overload [261]. Importantly, A $\beta$ -induced  $\text{Ca}^{2+}$  overload has not only been observed in neuronal cells but also in glial [262,447] and endothelial cells [448,449]. Especially glial cells have recently caught the attention of many leading AD research labs [450]. Atrophic and reactive astroglia were detected in AD mouse models [451], with reactive astrocytes being directly associated with A $\beta$  plaques [451] and NFTs [452]. Activated astrocytes and microglia furthermore modulate senile plaque formation, either promoting [453] or inhibiting plaque growth [454]. Elevated  $[\text{Ca}^{2+}]_i$  not only forms the basis for ER stress, glutathione depletion, ROS production and astrogliosis, but may furthermore stimulate the production and accumulation of A $\beta$  peptide in astrocytes [262,263]. Mutations in PS have been shown to sensitize astrocytes to ATP and glutamate [289], and A $\beta$  alters metabotropic  $\text{Ca}^{2+}$  signaling via a  $\text{Ca}^{2+}$ -calcineurin-nuclear factor of activated T-cells (NFAT)-dependent pathway. A $\beta$  peptide activates calpain, which in turn cleaves and activates calcineurin, a vital component of NFAT-mediated gene expression. Both calcineurin and calpain are well acknowledged as  $\text{Ca}^{2+}$  oscillation sensors [455,456] and NFAT-dependent gene expression is thus tightly linked to  $[\text{Ca}^{2+}]_i$  changes. Recent findings suggest that the  $\text{Ca}^{2+}$ -calcineurin-NFAT signaling axis is responsible for the overexpression of metabotropic GluR5, IP $_3$ R1 and IP $_3$ R2 in astrocytes surrounding A $\beta$  plaques, resulting in altered  $\text{Ca}^{2+}$  homeostasis [290,457]. Importantly, A $\beta$  differentially affects astrocyte metabotropic  $\text{Ca}^{2+}$  signaling in different brain regions which may account for the spatially distinct reactivity of astrocytes in the progression of AD [290].

Astrocyte ICWs have been shown to originate near A $\beta$  plaques and spread 200  $\mu\text{m}$  radially [21]. Based on the observation that photo-activation of caged IP $_3$  in astrocytes *in vivo* does not provoke ICWs, Crowe et al. [22] concluded that the spontaneous ICWs in AD mice do not depend on the IP $_3$ -signaling axis. Increases in Cx43 and Cx30 expression have been observed in reactive astrocytes in the immediate vicinity of the majority of A $\beta$  plaques in AD mouse models [313]. This is supported by an analysis performed in brain samples of AD patients, which displayed an increase in Cx43 expression [458]. A $\beta$  reduces GJIC and induces HC opening in cultured astrocytes and brain slices [367, 402], however, these observations are not supported by experimental results obtained from brain slices of APP-transgenic mice where GJ dye coupling through astrocytic networks was maintained in slices from different brain regions [459]. Interestingly, A $\beta$  peptide has been shown to lead to neuronal death via a cascade of HC activation, composed of both Cxs and Panxs. First, activated microglia promote the release of glutamate and ATP through microglial Cx43 HCs and Panx1 channels and astrocytic Cx43 HCs, which subsequently induce neuronal death by triggering Cx36 HC and Panx1 channel opening in neurons [402].

### 5.6.3. Traumatic brain and spinal cord injury

Traumatic brain and spinal cord injury is often complicated by delayed progressively expanding cell damage and death to which  $\text{Ca}^{2+}$  ions and Cxs can contribute [460,461]. Several studies have demonstrated a neuroprotective role of Cx channel blockers/Cx silencing in traumatic CNS injury [42,319,387,404,462,463]. Not only neuronal Cx36 GJs play a role in the secondary neuronal death [463,464], but also Cx43 channels can increase neuronal vulnerability to brain or spinal cord injury [318,319,387]. Astrocytic Cx43 furthermore plays a role in the development of chronic neuropathic pain following spinal cord injury [320]. Cx43 expression and phosphorylation are upregulated in response to traumatic CNS injury [308,309,321,465–467] and the phosphorylation of Cx43 is associated with autophagy in hippocampal pyramidal neurons [467]. In addition, Cx29 and Cx32 expression are *de novo* expressed in astrocytes and microglia localized in the injury border zone [468].

Astrocyte ICWs were observed *in vivo* in the border/penumbral region after traumatic injury. These ICWs were dependent on purinergic signaling, and affected the neuronal activity and survival in the border zone [24]. ATP release via astrocytic Cx43 HCs is suggested to aggravate the secondary injury [318]. Inositol polyphosphates are also released from stretch-injured astrocytes in culture, suggesting that release of IP $_3$  from glial cells could affect neighboring neurons [469]. In accordance, HCs have been put forth as IP $_3$ -releasing channels in the cochlea where the sensory neuroepithelium expresses IP $_3$ R on the plasma membrane [77]. It must be noted that IP $_3$ R expression has been identified in the plasma membrane of other cell types, but usually have the IP $_3$ -sensing domain on the intracellular site [470]. It thus remains an open question whether such signaling route can be activated from outside the cell.

### 5.6.4. Epilepsy

Excessive neuronal firing during epileptic seizures is associated with marked alterations in the composition of the extracellular ions, most notably an increase in  $\text{K}^+$  and a reduction of  $\text{Ca}^{2+}$  [471]. In accordance, lowering of  $[\text{Ca}^{2+}]_e$ , for example by stimulating glutamatergic synaptic transmission [254], has been shown to induce spontaneous epileptiform activity in hippocampal slices [472]. The resulting drop in extracellular  $\text{Ca}^{2+}$  stimulates CxHC opening that can further contribute to the translocation of  $\text{Ca}^{2+}$  ions to the intracellular compartment. Local or systemic administration of IP $_3$ -linked metabotropic glutamate receptor agonists can further contribute to excessive neuronal firing. Antagonists of these receptors indeed have anti-convulsant properties. Additionally, several anti-epileptic agents reduce the ability of astrocytes to transmit  $\text{Ca}^{2+}$  signaling [26].

Astrocytes isolated from human epilepsy patients display an increase in GJIC, and a surge in glutamate-induced  $\text{Ca}^{2+}$  oscillations and ICWs [473], which might be the consequence of an elevation in extracellular  $\text{K}^+$  concentration ( $[\text{K}^+]_e$ ) [474]. The contribution of Cx channels to ictogenesis is dual [4]. On the one hand, GJs may be anti-convulsive by contributing to spatial buffering of  $\text{K}^+$  and glutamate. The combined ablation of Cx43 and Cx30, which results in coupling-deficient astrocytes, lowers the threshold for epileptiform activity and thus supports this notion [52,57]. On the other hand, GJIC may be pro-convulsive by helping to provide neurons with the energy supply required for neuronal bursting, while ICWs propagated by GJs and CxHCs may expand the hyperexcited  $\text{Ca}^{2+}$  state of the astrocyte network thus modulating remotely located synapses [44,49]. In this respect, pharmacologic block of Cx channels reduced seizure activity while a potentiation of cell–cell coupling enhanced neuronal firing [52,388,475]. Interestingly, recent evidence based on work with a peptide targeting the second extracellular loop of Cx43 indicates that during an epileptiform insult, Cx43 HC opening is detrimental while GJIC is neuroprotective. By contrast, after the seizure, GJIC exacerbates neuronal death [388].

Astrocytes themselves may be the origin of glutamate in epileptic firing. Tian and co-workers [26] reported that paroxysmal depolarization shifts (PDSs, abnormally prolonged depolarizations with repetitive spiking manifesting as interictal discharges) can be initiated by the release of glutamate from extrasynaptic sources. The application of the  $\text{K}^+$  channel blocker 4-aminopyridine in the somatosensory cortex in vivo triggered PDSs, and stimulated the appearance of astrocytic ICWs and repetitive oscillatory  $\text{Ca}^{2+}$  increases. Astrocytic  $\text{Ca}^{2+}$  signaling invariably preceded bursting activity and flash photolysis of caged  $\text{Ca}^{2+}$  in astrocytes was sufficient to trigger PDSs. Additional support for an astrocytic origin is provided by the observation that intracellular dialysis of  $\text{IP}_3$  into single astrocytes evokes glutamate release and epileptiform discharges in adjacent neurons in hippocampal slices [476]. Two later studies demonstrated that glutamate release from astrocytes is not necessary to evoke epileptiform activity in hippocampal slices [477], but can evoke neuronal cell death in an in vivo model of status epilepticus [25].

## 6. Conclusions

For decades, research has been focusing on the neuronal abnormalities that accompany neurodegenerative disorders. However, at the same time, it is becoming increasingly evident that astrocytes are also important players in various CNS diseases. Reactive glial cells (mainly astrocytes and microglia) have been identified in many disorders of the CNS which boosted the interest in the potential role of these cells in neuronal injury. Astrocytes form extensive networks that connect neuronal synapses with the vasculature. This central position provides astrocytes with several essential functions, i.e. sensing neuronal activity as well as participating in their signaling, integrating electrochemical information and exchanging this information with the vasculature. Hereby, the canonical  $\text{PLC}/\text{IP}_3/\text{Ca}^{2+}$  pathway is the language via which astrocytes talk to each other as well as to neurons, microglia and vascular cells. The information processing within the glial network, as well as the communication of this information to NGVU partner cells is supported by intercellular signaling via GJs or via paracrine communication, possibly involving HCs. As such, it is no surprise that alterations of Cx channel function will profoundly affect proper functioning of the NGVU and will ultimately lead to cell dysfunction and cell death. An increasing number of publications have reported the occurrence of glial ICWs in vivo in various acute and chronic animal disease models such as epilepsy, ischemia, brain trauma and AD. The exact functional implications of these glial-based ICWs are unclear. As suggested here, they might provide the long-range transfer of information to evoke processes such as reactive astrogliosis, microglial activation and neurodegeneration. A balance between these processes likely determines the overall influence exerted by astrocytes in these conditions, saver versus killer,

and the level of destruction, protection and regeneration in neuropathological conditions. These examples of pathological conditions suggest further research to explore the terms and mechanisms leading to increased appearance of ICWs, and to determine their contribution in the pathogenesis and disease progression. Research on Cx-mediated  $\text{Ca}^{2+}$  communication within the NGVU may ultimately shed light on the development of targeted therapies for gliopathologies.

## Acknowledgements

This work is supported by the Fund for Scientific Research Flanders (FWO-Vlaanderen), Belgium (grant nos. G.0134.09, G.0298.11, G.0571.12 and G.0A54.13) and the Interuniversity Attraction Poles Program (Belgian Science Policy, projects P7/10 and P7/13). Figures were partly produced using Servier Medical art.

## References

- [1] A. Volterra, J. Meldolesi, Astrocytes, from brain glue to communication elements: the revolution continues, *Nat. Rev. Neurosci.* 6 (2005) 626–640.
- [2] C. Giaume, A. Koulakoff, L. Roux, D. Holcman, N. Rouach, Astroglial networks: a step further in neuroglial and gliovascular interactions, *Nat. Rev. Neurosci.* 11 (2010) 87–99.
- [3] N. Wang, M. De Bock, E. Decrock, M. Bol, A. Gadicherla, M. Vinken, V. Rogiers, F.F. Bukauskas, G. Bultynck, L. Leybaert, Paracrine signaling through plasma membrane hemichannels, *Biochim. Biophys. Acta* 1828 (2013) 35–50.
- [4] M. De Bock, M. Kerrebrouck, N. Wang, L. Leybaert, Neurological manifestations of oculodentodigital dysplasia: a Cx43 channelopathy of the central nervous system? *Front. Pharmacol.* 4 (2013) 120.
- [5] S. Penuela, R. Gehi, D.W. Laird, The biochemistry and function of pannexin channels, *Biochim. Biophys. Acta* 1828 (2013) 15–22.
- [6] A. Vogt, S.G. Hormuzdi, H. Monyer, Pannexin1 and Pannexin2 expression in the developing and mature rat brain, *Brain Res. Mol. Brain Res.* 141 (2005) 113–120.
- [7] J.I. Nagy, J.E. Rash, Connexins and gap junctions of astrocytes and oligodendrocytes in the CNS, *Brain Res. Brain Res. Rev.* 32 (2000) 29–44.
- [8] M. Theis, C. Giaume, Connexin-based intercellular communication and astrocyte heterogeneity, *Brain Res.* 1487 (2012) 88–98.
- [9] L. Leybaert, M.J. Sanderson, Intercellular  $\text{Ca}^{2+}$  waves: mechanisms and function, *Physiol. Rev.* 92 (2012) 1359–1392.
- [10] G. Dupont, L. Combettes, L. Leybaert, Calcium dynamics: spatio-temporal organization from the subcellular to the organ level, *Int. Rev. Cytol.* 261 (2007) 193–245.
- [11] E. Scemes, C. Giaume, Astrocyte calcium waves: what they are and what they do, *Glia* 54 (2006) 716–725.
- [12] G.J. Augustine, F. Santamaria, K. Tanaka, Local calcium signaling in neurons, *Neuron* 40 (2003) 331–346.
- [13] M.J. Berridge, Neuronal calcium signaling, *Neuron* 21 (1998) 13–26.
- [14] M.J. Berridge, Dysregulation of neural calcium signaling in Alzheimer disease, bipolar disorder and schizophrenia, *Prion* 7 (2013) 2–13.
- [15] A. Verkhratsky, J.J. Rodriguez, V. Parpura, Calcium signalling in astroglia, *Mol. Cell. Endocrinol.* 353 (2012) 45–56.
- [16] T.M. Hoogland, B. Kuhn, W. Gobel, W. Huang, J. Nakai, F. Helmchen, J. Flint, S.S. Wang, Radially expanding transglial calcium waves in the intact cerebellum, *Proc. Natl. Acad. Sci. U. S. A.* 106 (2009) 3496–3501.
- [17] A. Nimmerjahn, F. Kirchhoff, J.N. Kerr, F. Helmchen, Sulforhodamine 101 as a specific marker of astroglia in the neocortex in vivo, *Nat. Methods* 1 (2004) 31–37.
- [18] N. Kuga, T. Sasaki, Y. Takahara, N. Matsuki, Y. Ikegaya, Large-scale calcium waves traveling through astrocytic networks in vivo, *J. Neurosci.* 31 (2011) 2607–2614.
- [19] C. Mathiesen, A. Brazhe, K. Thomsen, M. Lauritzen, Spontaneous calcium waves in Bergman glia increase with age and hypoxia and may reduce tissue oxygen, *J. Cereb. Blood Flow Metab.* 33 (2013) 161–169.
- [20] S. Ding, T. Wang, W. Cui, P.G. Haydon, Photothrombosis ischemia stimulates a sustained astrocytic  $\text{Ca}^{2+}$  signaling in vivo, *Glia* 57 (2009) 767–776.
- [21] K.V. Kuchibhotla, C.R. Lattarulo, B.T. Hyman, B.J. Bacskai, Synchronous hyperactivity and intercellular calcium waves in astrocytes in Alzheimer mice, *Science* 323 (2009) 1211–1215.
- [22] S.E. Crowe, S. Kantevari, G.C. Ellis-Davies, Photochemically initiated intracellular astrocytic calcium waves in living mice using two-photon uncaging of  $\text{IP}_3$ , *ACS Chem. Neurosci.* 1 (2010) 575–585.
- [23] D. Sieger, C. Moritz, T. Ziegenhals, S. Prykhodzhiy, F. Peri, Long-range  $\text{Ca}^{2+}$  waves transmit brain-damage signals to microglia, *Dev. Cell* 22 (2012) 1138–1148.
- [24] A.M. Choo, W.J. Miller, Y.C. Chen, P. Nibley, T.P. Patel, C. Goletiani, B. Morrison III, M. K. Kutzing, B.L. Firestein, J.Y. Sul, P.G. Haydon, D.F. Meaney, Antagonism of purinergic signalling improves recovery from traumatic brain injury, *Brain* 136 (2013) 65–80.
- [25] S. Ding, T. Fellin, Y. Zhu, S.Y. Lee, Y.P. Auberson, D.F. Meaney, D.A. Coulter, G. Carmignoto, P.G. Haydon, Enhanced astrocytic  $\text{Ca}^{2+}$  signals contribute to neuronal excitotoxicity after status epilepticus, *J. Neurosci.* 27 (2007) 10674–10684.
- [26] G.F. Tian, H. Azmi, T. Takano, Q. Xu, W. Peng, J. Lin, N. Oberheim, N. Lou, X. Wang, H. R. Zielke, J. Kang, M. Nedergaard, An astrocytic basis of epilepsy, *Nat. Med.* 11 (2005) 973–981.

- [27] E.A. Neuwelt, B. Bauer, C. Fahlke, G. Fricker, C. Iadecola, D. Janigro, L. Leybaert, Z. Molnar, M.E. O'Donnell, J.T. Povlishock, N.R. Saunders, F. Sharp, D. Stanimirovic, R.J. Watts, L.R. Drewes, Engaging neuroscience to advance translational research in brain barrier biology, *Nat. Rev. Neurosci.* 12 (2011) 169–182.
- [28] B.T. Hawkins, T.P. Davis, The blood–brain barrier/neurovascular unit in health and disease, *Pharmacol. Rev.* 57 (2005) 173–185.
- [29] E.A. Bushong, M.E. Martone, Y.Z. Jones, M.H. Ellisman, Protoplasmic astrocytes in CA1 stratum radiatum occupy separate anatomical domains, *J. Neurosci.* 22 (2002) 183–192.
- [30] N.A. Oberheim, X. Wang, S. Goldman, M. Nedergaard, Astrocytic complexity distinguishes the human brain, *Trends Neurosci.* 29 (2006) 547–553.
- [31] V. Parpura, T.A. Basarsky, F. Liu, K. Jeftinija, S. Jeftinija, P.G. Haydon, Glutamate-mediated astrocyte–neuron signalling, *Nature* 369 (1994) 744–747.
- [32] E.A. Newman, Calcium signaling in retinal glial cells and its effect on neuronal activity, *Prog. Brain Res.* 132 (2001) 241–254.
- [33] G. Perea, A. Araque, GLIA modulates synaptic transmission, *Brain Res. Rev.* 63 (2010) 93–102.
- [34] T.A. Fiacco, K.D. McCarthy, Astrocyte calcium elevations: properties, propagation, and effects on brain signaling, *Glia* 54 (2006) 676–690.
- [35] E. Shigetomi, O. Jackson-Weaver, R.T. Huckstepp, T.J. O'Dell, B.S. Khakh, TRPA1 channels are regulators of astrocyte basal calcium levels and long-term potentiation via constitutive D-serine release, *J. Neurosci.* 33 (2013) 10143–10153.
- [36] M. Santello, C. Cali, P. Bezzi, Gliotransmission and the tripartite synapse, *Adv. Exp. Med. Biol.* 970 (2012) 307–331.
- [37] J.T. Porter, K.D. McCarthy, Astrocytic neurotransmitter receptors *in situ* and *in vivo*, *Prog. Neurobiol.* 51 (1997) 439–455.
- [38] J.A. Filosa, A.D. Bonev, M.T. Nelson, Calcium dynamics in cortical astrocytes and arterioles during neurovascular coupling, *Circ. Res.* 95 (2004) e73–e81.
- [39] C. Iadecola, M. Nedergaard, Glial regulation of the cerebral microvasculature, *Nat. Neurosci.* 10 (2007) 1369–1376.
- [40] R.C. Koehler, D. Gebremedhin, D.R. Harder, Role of astrocytes in cerebrovascular regulation, *J. Appl. Physiol.* 100 (2006) 307–317.
- [41] B. Nadarajah, J.G. Parnavelas, Gap junction-mediated communication in the developing and adult cerebral cortex, *Novartis Found. Symp.* 219 (1999) 157–170 (discussion 170–154).
- [42] A.B. Belousov, J.D. Fontes, Neuronal gap junctions: making and breaking connections during development and injury, *Trends Neurosci.* 36 (2013) 227–236.
- [43] C. Cina, J.F. Bechberger, M.A. Ozog, C.C. Naus, Expression of connexins in embryonic mouse neocortical development, *J. Comp. Neurol.* 504 (2007) 298–313.
- [44] C. Giaume, Astroglial wiring is adding complexity to neuroglial networking, *Front. Neuroenerg.* 2 (2010).
- [45] C. Giaume, M. Theis, Pharmacological and genetic approaches to study connexin-mediated channels in glial cells of the central nervous system, *Brain Res. Rev.* 63 (2010) 160–176.
- [46] F. Blomstrand, N.D. Aberg, P.S. Eriksson, E. Hansson, L. Ronnback, Extent of intercellular calcium wave propagation is related to gap junction permeability and level of connexin-43 expression in astrocytes in primary cultures from four brain regions, *Neuroscience* 92 (1999) 255–265.
- [47] R. Dermietzel, E.L. Hertberg, J.A. Kessler, D.C. Spray, Gap junctions between cultured astrocytes: immunocytochemical, molecular, and electrophysiological analysis, *J. Neurosci.* 11 (1991) 1421–1432.
- [48] P. Ezan, P. Andre, S. Cisternino, B. Saubamea, A.C. Boulay, S. Dautremer, M.A. Thomas, N. Quenec'h du, C. Giaume, M. Cohen-Salmon, Deletion of astroglial connexins weakens the blood–brain barrier, *J. Cereb. Blood Flow Metab.* 32 (2012) 1457–1467.
- [49] N. Rouach, A. Koulakoff, V. Abudara, K. Willecke, C. Giaume, Astroglial metabolic networks sustain hippocampal synaptic transmission, *Science* 322 (2008) 1551–1555.
- [50] M. Simard, G. Arcuino, T. Takano, Q.S. Liu, M. Nedergaard, Signaling at the gliovascular interface, *J. Neurosci.* 23 (2003) 9254–9262.
- [51] A. Koulakoff, P. Ezan, C. Giaume, Neurons control the expression of connexin 30 and connexin 43 in mouse cortical astrocytes, *Glia* 56 (2008) 1299–1311.
- [52] A. Wallraff, R. Kohling, U. Heinemann, M. Theis, K. Willecke, C. Steinhauser, The impact of astrocytic gap junctional coupling on potassium buffering in the hippocampus, *J. Neurosci.* 26 (2006) 5438–5447.
- [53] J.I. Nagy, B.D. Lynn, O. Tress, K. Willecke, J.E. Rash, Connexin26 expression in brain parenchymal cells demonstrated by targeted connexin ablation in transgenic mice, *Eur. J. Neurosci.* 34 (2011) 263–271.
- [54] R. Dermietzel, Y. Gao, E. Scemes, D. Vieira, M. Urban, M. Kremer, M.V. Bennett, D.C. Spray, Connexin43 null mice reveal that astrocytes express multiple connexins, *Brain Res. Brain Res. Rev.* 32 (2000) 45–56.
- [55] E. Scemes, R. Dermietzel, D.C. Spray, Calcium waves between astrocytes from Cx43 knockout mice, *Glia* 24 (1998) 65–73.
- [56] N. Rouach, J. Glowinski, C. Giaume, Activity-dependent neuronal control of gap-junctional communication in astrocytes, *J. Cell Biol.* 149 (2000) 1513–1526.
- [57] M. Theis, R. Jauch, L. Zhuo, D. Speidel, A. Wallraff, B. Doring, C. Frisch, G. Sohl, B. Teubner, C. Euwens, J. Huston, C. Steinhauser, A. Messing, U. Heinemann, K. Willecke, Accelerated hippocampal spreading depression and enhanced locomotor activity in mice with astrocyte-directed inactivation of connexin43, *J. Neurosci.* 23 (2003) 766–776.
- [58] H. Marrero, R.K. Orkand, Nerve impulses increase glial intercellular permeability, *Glia* 16 (1996) 285–289.
- [59] H.L. Xu, L. Mao, S. Ye, C. Paisansathan, F. Vetri, D.A. Pelligrino, Astrocytes are a key conduit for upstream signaling of vasodilation during cerebral cortical neuronal activation *in vivo*, *Am. J. Physiol. Heart Circ. Physiol.* 294 (2008) H622–H632.
- [60] J.E. Rash, T. Yasumura, K.G. Davidson, C.S. Furman, F.E. Dudek, J.I. Nagy, Identification of cells expressing Cx43, Cx30, Cx26, Cx32 and Cx36 in gap junctions of rat brain and spinal cord, *Cell Commun. Adhes.* 8 (2001) 315–320.
- [61] M. Nedergaard, Direct signaling from astrocytes to neurons in cultures of mammalian brain cells, *Science* 263 (1994) 1768–1771.
- [62] P. Pakhotin, A. Verkhratsky, Electrical synapses between Bergmann glial cells and Purkinje neurones in rat cerebellar slices, *Mol. Cell. Neurosci.* 28 (2005) 79–84.
- [63] L. Leybaert, K. Paemeleire, A. Strahonja, M.J. Sanderson, Inositol-trisphosphate-dependent intercellular calcium signaling in and between astrocytes and endothelial cells, *Glia* 24 (1998) 398–407.
- [64] M.M. Froes, A.H. Correia, J. Garcia-Abreu, D.C. Spray, A.C. Campos de Carvalho, M.V. Neto, Gap-junctional coupling between neurons and astrocytes in primary central nervous system cultures, *Proc. Natl. Acad. Sci. U. S. A.* 96 (1999) 7541–7546.
- [65] K. Bittman, D.L. Becker, F. Cicirata, J.G. Parnavelas, Connexin expression in homotypic and heterotypic cell coupling in the developing cerebral cortex, *J. Comp. Neurol.* 443 (2002) 201–212.
- [66] L. Duan, H. Yuan, C.J. Su, Y.Y. Liu, Z.R. Rao, Ultrastructure of junction areas between neurons and astrocytes in rat supraoptic nuclei, *World J. Gastroenterol.* 10 (2004) 117–121.
- [67] V. Alvarez-Maubecin, F. Garcia-Hernandez, J.T. Williams, E.J. Van Bockstaele, Functional coupling between neurons and glia, *J. Neurosci.* 20 (2000) 4091–4098.
- [68] K. Braet, K. Paemeleire, K. D'Herde, M.J. Sanderson, L. Leybaert, Astrocyte-endothelial cell calcium signals conveyed by two signalling pathways, *Eur. J. Neurosci.* 13 (2001) 79–91.
- [69] C. Giaume, L. Leybaert, C.C. Naus, J.C. Saez, Connexin and pannexin hemichannels in brain glial cells: properties, pharmacology, and roles, *Front. Pharmacol.* 4 (2013) 88.
- [70] M.V. Bennett, J.E. Contreras, F.F. Bukauskas, J.C. Saez, New roles for astrocytes: gap junction hemichannels have something to communicate, *Trends Neurosci.* 26 (2003) 610–617.
- [71] W.H. Evans, E. De Vuyst, L. Leybaert, The gap junction cellular internet: connexin hemichannels enter the signalling limelight, *Biochem. J.* 397 (2006) 1–14.
- [72] J. Kang, N. Kang, D. Lovatt, A. Torres, Z. Zhao, J. Lin, M. Nedergaard, Connexin 43 hemichannels are permeable to ATP, *J. Neurosci.* 28 (2008) 4702–4711.
- [73] Z.C. Ye, M.S. Wyeth, S. Baltan-Tekkuk, B.R. Ransom, Functional hemichannels in astrocytes: a novel mechanism of glutamate release, *J. Neurosci.* 23 (2003) 3588–3596.
- [74] S. Burra, J.X. Jiang, Connexin 43 hemichannel opening associated with Prostaglandin E(2) release is adaptively regulated by mechanical stimulation, *Commun. Integr. Biol.* 2 (2009) 239–240.
- [75] H. Okuda, K. Nishida, Y. Higashi, K. Nagasawa, NAD(+) influx through connexin hemichannels prevents poly(ADP-ribose) polymerase-mediated astrocyte death, *Life Sci.* 92 (2013) 808–814.
- [76] S. Bruzzone, L. Guida, E. Zocchi, L. Franco, A. De Flora, Connexin 43 hemi channels mediate Ca<sup>2+</sup> + -regulated transmembrane NAD<sup>+</sup> fluxes in intact cells, *FASEB J.* 15 (2001) 10–12.
- [77] D.G. Gossman, H.B. Zhao, Hemichannel-mediated inositol 1,4,5-trisphosphate (IP<sub>3</sub>) release in the cochlea: a novel mechanism of IP<sub>3</sub> intercellular signaling, *Cell Commun. Adhes.* 15 (2008) 305–315.
- [78] M.A. Retamal, N. Froger, N. Palacios-Prado, P. Ezan, P.J. Saez, J.C. Saez, C. Giaume, Cx43 hemichannels and gap junction channels in astrocytes are regulated oppositely by proinflammatory cytokines released from activated microglia, *J. Neurosci.* 27 (2007) 13781–13792.
- [79] J.E. Contreras, J.C. Saez, F.F. Bukauskas, M.V. Bennett, Gating and regulation of connexin 43 (Cx43) hemichannels, *Proc. Natl. Acad. Sci. U. S. A.* 100 (2003) 11388–11393.
- [80] J.C. Saez, M.A. Retamal, D. Basilio, F.F. Bukauskas, M.V. Bennett, Connexin-based gap junction hemichannels: gating mechanisms, *Biochim. Biophys. Acta* 1711 (2005) 215–224.
- [81] D.C. Spray, Z.C. Ye, B.R. Ransom, Functional connexin “hemichannels”: a critical appraisal, *Glia* 54 (2006) 758–773.
- [82] J.A. Orellana, E. Diaz, K.A. Schalper, A.A. Vargas, M.V. Bennett, J.C. Saez, Cation permeation through connexin 43 hemichannels is cooperative, competitive and saturable with parameters depending on the permeant species, *Biochem. Biophys. Res. Commun.* 409 (2011) 603–609.
- [83] E. De Vuyst, E. Decrock, M. De Bock, H. Yamasaki, C.C. Naus, W.H. Evans, L. Leybaert, Connexin hemichannels and gap junction channels are differentially influenced by lipopolysaccharide and basic fibroblast growth factor, *Mol. Biol. Cell* 18 (2007) 34–46.
- [84] J.E. Contreras, H.A. Sanchez, E.A. Eugenin, D. Speidel, M. Theis, K. Willecke, F.F. Bukauskas, M.V. Bennett, J.C. Saez, Metabolic inhibition induces opening of unapposed connexin 43 gap junction hemichannels and reduces gap junctional communication in cortical astrocytes in culture, *Proc. Natl. Acad. Sci. U. S. A.* 99 (2002) 495–500.
- [85] S. Ramachandran, L.H. Xie, S.A. John, S. Subramaniam, R. Lal, A novel role for connexin hemichannel in oxidative stress and smoking-induced cell injury, *PLoS One* 2 (2007) e712.
- [86] E. De Vuyst, N. Wang, E. Decrock, M. De Bock, M. Vinken, M. Van Moorhem, C. Lai, M. Culet, V. Rogiers, R. Cecchetti, C.C. Naus, W.H. Evans, L. Leybaert, Ca(2+) regulation of connexin 43 hemichannels in C6 glioma and glial cells, *Cell Calcium* 46 (2009) 176–187.
- [87] J. Thimm, A. Mechler, H. Lin, S. Rhee, R. Lal, Calcium-dependent open/closed conformations and interfacial energy maps of reconstituted hemichannels, *J. Biol. Chem.* 280 (2005) 10646–10654.
- [88] A. Chandrasekhar, A.K. Bera, Hemichannels: permeants and their effect on development, physiology and death, *Cell Biochem. Funct.* 30 (2012) 89–100.



- [89] E. De Vuyst, E. Decrock, L. Cabooter, G.R. Dubyak, C.C. Naus, W.H. Evans, L. Leybaert, Intracellular calcium changes trigger connexin 32 hemichannel opening, *EMBO J.* 25 (2006) 34–44.
- [90] D.J. Belliveau, M. Bani-Yaghoob, B. McGirr, C.C. Naus, W.J. Rushlow, Enhanced neurite outgrowth in PC12 cells mediated by connexin hemichannels and ATP, *J. Biol. Chem.* 281 (2006) 20920–20931.
- [91] K.A. Schalter, N. Palacios-Prado, M.A. Retamal, K.F. Shoji, A.D. Martinez, J.C. Saez, Connexin hemichannel composition determines the FGF-1-induced membrane permeability and free  $[Ca^{2+}]_i$  responses, *Mol. Biol. Cell* 19 (2008) 3501–3513.
- [92] J. Stehberg, R. Moraga-Amaro, C. Salazar, A. Becerra, C. Echeverria, J.A. Orellana, G. Bultynck, R. Ponsaerts, L. Leybaert, F. Simon, J.C. Saez, M.A. Retamal, Release of gliotransmitters through astroglial connexin 43 hemichannels is necessary for fear memory consolidation in the basolateral amygdala, *FASEB J.* 26 (2011) 3649–3657.
- [93] R. Iglesias, G. Dahl, F. Qiu, D.C. Spray, E. Scemes, Pannexin 1: the molecular substrate of astrocyte “hemichannels”, *J. Neurosci.* 29 (2009) 7092–7097.
- [94] S.O. Suadicani, R. Iglesias, J. Wang, G. Dahl, D.C. Spray, E. Scemes, ATP signaling is deficient in cultured Pannexin1-null mouse astrocytes, *Glia* 60 (2012) 1106–1116.
- [95] Y. Huang, J.B. Grinspan, C.K. Abrams, S.S. Scherer, Pannexin1 is expressed by neurons and glia but does not form functional gap junctions, *Glia* 55 (2007) 46–56.
- [96] G.E. Sosinsky, D. Boassa, R. Dermietzel, H.S. Duffy, D.W. Laird, B. MacVicar, C.C. Naus, S. Penuela, E. Scemes, D.C. Spray, R.J. Thompson, H.B. Zhao, G. Dahl, Pannexin channels are not gap junction hemichannels, *Channels (Austin)* 5 (2011) 193–197.
- [97] J.X. Jiang, S. Gu, Gap junction- and hemichannel-independent actions of connexins, *Biochim. Biophys. Acta* 1711 (2005) 208–214.
- [98] M. Vinken, E. Decrock, E. De Vuyst, R. Ponsaerts, C. D’Hondt, G. Bultynck, L. Ceelen, T. Vanhaecke, L. Leybaert, V. Rogiers, Connexins: sensors and regulators of cell cycling, *Biochim. Biophys. Acta* 1815 (2011) 13–25.
- [99] D.A. Jacobas, E. Scemes, D.C. Spray, Gene expression alterations in connexin null mice extend beyond the gap junction, *Neurochem. Int.* 45 (2004) 243–250.
- [100] D.A. Jacobas, S. Jacobas, D.C. Spray, Connexin-dependent transcriptomic networks in mouse brain, *Prog. Biophys. Mol. Biol.* 94 (2007) 169–185.
- [101] E. Scemes, Modulation of astrocyte P2Y1 receptors by the carboxyl terminal domain of the gap junction protein Cx43, *Glia* 56 (2008) 145–153.
- [102] E. Scemes, S.O. Suadicani, D.C. Spray, Intercellular communication in spinal cord astrocytes: fine tuning between gap junctions and P2 nucleotide receptors in calcium wave propagation, *J. Neurosci.* 20 (2000) 1435–1445.
- [103] S. Oik, A. Turchinovich, M. Grzondowski, K. Stuhler, H.E. Meyer, G. Zoidl, R. Dermietzel, Proteomic analysis of astroglial connexin43 silencing uncovers a cytoskeletal platform involved in process formation and migration, *Glia* 58 (2010) 494–505.
- [104] L.A. Elias, D.D. Wang, A.R. Kriegstein, Gap junction adhesion is necessary for radial migration in the neocortex, *Nature* 448 (2007) 901–907.
- [105] C. Cina, K. Maass, M. Theis, K. Willecke, J.F. Bechberger, C.C. Naus, Involvement of the cytoplasmic C-terminal domain of connexin43 in neuronal migration, *J. Neurosci.* 29 (2009) 2009–2021.
- [106] M.J. Berridge, P. Lipp, M.D. Bootman, The versatility and universality of calcium signalling, *Nat. Rev. Mol. Cell Biol.* 1 (2000) 11–21.
- [107] J.R. Lopez, A. Lyckman, S. Oddo, F.M. Laferla, H.W. Querfurth, A. Shtifman, Increased intraneuronal resting  $[Ca^{2+}]_i$  in adult Alzheimer’s disease mice, *J. Neurochem.* 105 (2008) 262–271.
- [108] R.M. Case, D. Eisner, A. Gurney, O. Jones, S. Muallem, A. Verkhratsky, Evolution of calcium homeostasis: from birth of the first cell to an omnipresent signalling system, *Cell Calcium* 42 (2007) 345–350.
- [109] G.G. Somjen, The regulation of brain ions, *Ions in the brain: normal function, seizures and stroke*, Oxford University Press, 2004, pp. 13–62.
- [110] M. Massimini, F. Amzica, Extracellular calcium fluctuations and intracellular potentials in the cortex during the slow sleep oscillation, *J. Neurophysiol.* 85 (2001) 1346–1350.
- [111] G.G. Somjen, Ion regulation in the brain: implications for pathophysiology, *Neuroscientist* 8 (2002) 254–267.
- [112] J. Meldolesi, T. Pozzan, The endoplasmic reticulum  $Ca^{2+}$  store: a view from the lumen, *Trends Biochem. Sci.* 23 (1998) 10–14.
- [113] J.Y. Chatton, H. Liu, J.W. Stucki, Simultaneous measurements of  $Ca^{2+}$  in the intracellular stores and the cytosol of hepatocytes during hormone-induced  $Ca^{2+}$  oscillations, *FEBS Lett.* 368 (1995) 165–168.
- [114] P.G. Wood, J.I. Gillespie, Evidence for mitochondrial  $Ca^{2+}$ -induced  $Ca^{2+}$  release in permeabilised endothelial cells, *Biochem. Biophys. Res. Commun.* 246 (1998) 543–548.
- [115] R. Rizzuto, D. De Stefani, A. Raffaello, C. Mammucari, Mitochondria as sensors and regulators of calcium signalling, *Nat. Rev. Mol. Cell Biol.* 13 (2012) 566–578.
- [116] M.J. Berridge, Inositol trisphosphate and calcium signalling mechanisms, *Biochim. Biophys. Acta* 1793 (2009) 933–940.
- [117] A. Verkhratsky, V. Pappas, Store-operated calcium entry in neuroglia, *Neurosci. Bull.* 30 (2014) 125–133.
- [118] J.W. Putney, Alternative forms of the store-operated calcium entry mediators, STIM1 and Orai1, *Curr. Top. Membr.* 71 (2013) 109–123.
- [119] J.W. Putney, Capacitative calcium entry: from concept to molecules, *Immunol. Rev.* 231 (2009) 10–22.
- [120] J.H. Hong, Q. Li, M.S. Kim, D.M. Shin, S. Feske, L. Birnbaumer, K.T. Cheng, I.S. Ambudkar, S. Muallem, Polarized but differential localization and recruitment of STIM1, Orai1 and TRPC channels in secretory cells, *Traffic* 12 (2011) 232–245.
- [121] G.M. Salido, I. Jardin, J.A. Rosado, The TRPC ion channels: association with Orai1 and STIM1 proteins and participation in capacitative and non-capacitative calcium entry, *Adv. Exp. Med. Biol.* 704 (2011) 413–433.
- [122] E. Samuels, J.B. Parys, L. Missiaen, H. De Smedt, G. Bultynck, Intracellular  $Ca^{2+}$  storage in health and disease: a dynamic equilibrium, *Cell Calcium* 47 (2010) 297–314.
- [123] J.B. Parys, H. De Smedt, Inositol 1,4,5-trisphosphate and its receptors, *Adv. Exp. Med. Biol.* 740 (2012) 255–279.
- [124] K. Mikoshiba, IP3 receptor/ $Ca^{2+}$  channel: from discovery to new signaling concepts, *J. Neurochem.* 102 (2007) 1426–1446.
- [125] J.T. Lanner, Ryanodine receptor physiology and its role in disease, *Adv. Exp. Med. Biol.* 740 (2012) 217–234.
- [126] J.K. Foskett, C. White, K.H. Cheung, D.O. Mak, Inositol trisphosphate receptor  $Ca^{2+}$  release channels, *Physiol. Rev.* 87 (2007) 593–658.
- [127] H.L. Roderick, M.J. Berridge, M.D. Bootman, Calcium-induced calcium release, *Curr. Biol.* 13 (2003) R425.
- [128] C.M. Vines, Phospholipase C, *Adv. Exp. Med. Biol.* 740 (2012) 235–254.
- [129] S.V. Straub, A.D. Bonev, M.K. Wilkerson, M.T. Nelson, Dynamic inositol trisphosphate-mediated calcium signals within astrocytic endfeet underlie vasodilation of cerebral arterioles, *J. Gen. Physiol.* 128 (2006) 659–669.
- [130] I. Parker, J. Choi, Y. Yao, Elementary events of InsP3-induced  $Ca^{2+}$  liberation in *Xenopus* oocytes: hot spots, puffs and blips, *Cell Calcium* 20 (1996) 105–121.
- [131] T. Cali, D. Ottoloni, M. Brini, Mitochondrial  $Ca^{2+}$  as a key regulator of mitochondrial activities, *Adv. Exp. Med. Biol.* 942 (2012) 53–73.
- [132] V. Shoshan-Barmatz, M. Golan, Mitochondrial VDAC1: function in cell life and death and a target for cancer therapy, *Curr. Med. Chem.* 19 (2012) 714–735.
- [133] F. Perocchi, V.M. Gohil, H.S. Garg, X.R. Bao, J.E. McCombs, A.E. Palmer, V.K. Mootha, MICU1 encodes a mitochondrial EF hand protein required for  $Ca^{2+}$  uptake, *Nature* 467 (2010) 291–296.
- [134] J.M. Baughman, F. Perocchi, H.S. Garg, M. Plovanich, C.A. Belcher-Timme, Y. Sancak, X.R. Bao, L. Strittmatter, O. Goldberg, R.L. Bogorad, V. Kotliarsky, V.K. Mootha, Integrative genomics identifies MCU as an essential component of the mitochondrial calcium uniporter, *Nature* 476 (2011) 341–345.
- [135] D. De Stefani, A. Raffaello, E. Teardo, I. Szabo, R. Rizzuto, A forty-kilodalton protein of the inner membrane is the mitochondrial calcium uniporter, *Nature* 476 (2011) 336–340.
- [136] K. Mallilankaraman, C. Cardenas, P.J. Doonan, H.C. Chandramoorthy, K.M. Irrinki, T. Golenar, G. Scordas, P. Madireddi, J. Yang, M. Muller, R. Miller, J.E. Kolesar, J. Molgo, B. Kaufman, G. Hajnoczky, J.K. Foskett, M. Madesh, MCU1 is an essential component of mitochondrial  $Ca^{2+}$  uptake that regulates cellular metabolism, *Nat. Cell Biol.* 14 (2012) 1336–1343.
- [137] R. Rizzuto, M. Brini, M. Murgia, T. Pozzan, Microdomains with high  $Ca^{2+}$  close to IP3-sensitive channels that are sensed by neighboring mitochondria, *Science* 262 (1993) 744–747.
- [138] G. Scordas, A.P. Thomas, G. Hajnoczky, Quasi-synaptic calcium signal transmission between endoplasmic reticulum and mitochondria, *EMBO J.* 18 (1999) 96–108.
- [139] G. Scordas, C. Renken, P. Varnai, L. Walter, D. Weaver, K.F. Buttler, T. Balla, C.A. Mannella, G. Hajnoczky, Structural and functional features and significance of the physical linkage between ER and mitochondria, *J. Cell Biol.* 174 (2006) 915–921.
- [140] R. Rizzuto, P. Pinton, W. Carrington, F.S. Fay, K.E. Fogarty, L.M. Lifshitz, R.A. Tuft, T. Pozzan, Close contacts with the endoplasmic reticulum as determinants of mitochondrial  $Ca^{2+}$  responses, *Science* 280 (1998) 1763–1766.
- [141] G. Scordas, P. Varnai, T. Golenar, S. Roy, G. Purkins, T.G. Schneider, T. Balla, G. Hajnoczky, Imaging interorganelle contacts and local calcium dynamics at the ER-mitochondrial interface, *Mol. Cell* 39 (2010) 121–132.
- [142] A.E. Rusinol, Z. Cui, M.H. Chen, J.E. Vance, A unique mitochondria-associated membrane fraction from rat liver has a high capacity for lipid synthesis and contains pre-Golgi secretory proteins including nascent lipoproteins, *J. Biol. Chem.* 269 (1994) 27494–27502.
- [143] S. Marchi, M. Marinello, A. Bononi, M. Bonora, C. Giorgi, A. Rimessi, P. Pinton, Selective modulation of subtype III IP(3)R by Akt regulates ER  $Ca^{2+}$  release and apoptosis, *Cell Death Dis.* 3 (2012) e304.
- [144] C. Bionda, J. Portoukalian, D. Schmitt, C. Rodriguez-Lafrasse, D. Ardail, Subcellular compartmentalization of ceramide metabolism: MAM (mitochondria-associated membrane) and/or mitochondria? *Biochem. J.* 382 (2004) 527–533.
- [145] D. Ardail, I. Popa, J. Bodenne, P. Louisot, D. Schmitt, J. Portoukalian, The mitochondria-associated endoplasmic-reticulum subcompartment (MAM fraction) of rat liver contains highly active sphingolipid-specific glycosyltransferases, *Biochem. J.* 371 (2003) 1013–1019.
- [146] A. Bononi, S. Missiroli, F. Poletti, J.M. Suski, C. Agnoletto, M. Bonora, E. De Marchi, C. Giorgi, S. Marchi, S. Patergnani, A. Rimessi, M.R. Wieckowski, P. Pinton, Mitochondria-associated membranes (MAMs) as hotspot  $Ca^{2+}$  signaling units, *Adv. Exp. Med. Biol.* 740 (2012) 411–437.
- [147] O.M. de Brito, L. Scorrano, An intimate liaison: spatial organization of the endoplasmic reticulum-mitochondria relationship, *EMBO J.* 29 (2010) 2715–2723.
- [148] A. Raturi, T. Simmen, Where the endoplasmic reticulum and the mitochondrion tie the knot: the mitochondria-associated membrane (MAM), *Biochim. Biophys. Acta* 1833 (2013) 213–224.
- [149] C. Giorgi, D. De Stefani, A. Bononi, R. Rizzuto, P. Pinton, Structural and functional link between the mitochondrial network and the endoplasmic reticulum, *Int. J. Biochem. Cell Biol.* 41 (2009) 1817–1827.
- [150] E.A. Schon, E. Area-Gomez, Mitochondria-associated ER membranes in Alzheimer disease, *Mol. Cell. Neurosci.* 55 (2013) 26–36.
- [151] G. Szabadkai, K. Bianchi, P. Varnai, D. De Stefani, M.R. Wieckowski, D. Cavagna, A.I. Nagy, T. Balla, R. Rizzuto, Chaperone-mediated coupling of endoplasmic reticulum and mitochondrial  $Ca^{2+}$  channels, *J. Cell Biol.* 175 (2006) 901–911.
- [152] O.M. de Brito, L. Scorrano, Mitofusin 2 tethers endoplasmic reticulum to mitochondria, *Nature* 456 (2008) 605–610.



- [153] T. Simmen, J.E. Aslan, A.D. Blagoveshchenskaya, L. Thomas, L. Wan, Y. Xiang, S.F. Feliciangeli, C.H. Hung, C.M. Crump, G. Thomas, PACS-2 controls endoplasmic reticulum-mitochondria communication and Bid-mediated apoptosis, *EMBO J.* 24 (2005) 717–729.
- [154] T.J. Collins, P. Lipp, M.J. Berridge, M.D. Bootman, Mitochondrial  $\text{Ca}^{2+}$  uptake depends on the spatial and temporal profile of cytosolic  $\text{Ca}^{2+}$  signals, *J. Biol. Chem.* 276 (2001) 26411–26420.
- [155] L.S. Jouaville, P. Pinton, C. Bastianutto, G.A. Rutter, R. Rizzuto, Regulation of mitochondrial ATP synthesis by calcium: evidence for a long-term metabolic priming, *Proc. Natl. Acad. Sci. U. S. A.* 96 (1999) 13807–13812.
- [156] C. Cardenas, R.A. Miller, I. Smith, T. Bui, J. Molgo, M. Muller, H. Vais, K.H. Cheung, J. Yang, I. Parker, C.B. Thompson, M.J. Birnbaum, K.R. Hallows, J.K. Foskett, Essential regulation of cell bioenergetics by constitutive  $\text{InsP}_3$  receptor  $\text{Ca}^{2+}$  transfer to mitochondria, *Cell* 142 (2010) 270–283.
- [157] M. Bonora, S. Patergnani, A. Rimesi, E. De Marchi, J.M. Suski, A. Bononi, C. Giorgi, S. Marchi, S. Missiroli, F. Poletti, M.R. Wieckowski, P. Pinton, ATP synthesis and storage, *Purinergic Signal* 8 (2012) 343–357.
- [158] R. Palty, W.F. Silverman, M. Hershfinkel, T. Caporale, S.L. Sensi, J. Parnis, C. Nolte, D. Fishman, V. Shoshan-Barmatz, S. Herrmann, D. Khananshvil, I. Sekler, NCLX is an essential component of mitochondrial  $\text{Na}^+/\text{Ca}^{2+}$  exchange, *Proc. Natl. Acad. Sci. U. S. A.* 107 (2010) 436–441.
- [159] R. Palty, I. Sekler, The mitochondrial  $\text{Na}^+/\text{Ca}^{2+}$  exchanger, *Cell Calcium* 52 (2012) 9–15.
- [160] T. Cali, D. Ottoloni, M. Brini, Calcium and endoplasmic reticulum-mitochondria tethering in neurodegeneration, *DNA Cell Biol.* 32 (2013) 140–146.
- [161] N.L. Allbritton, T. Meyer, L. Stryer, Range of messenger action of calcium ion and inositol 1,4,5-trisphosphate, *Science* 258 (1992) 1812–1815.
- [162] G.C. Churchill, C.F. Louis, Roles of  $\text{Ca}^{2+}$ , inositol trisphosphate and cyclic ADP-ribose in mediating intercellular  $\text{Ca}^{2+}$  signaling in sheep lens cells, *J. Cell Sci.* 111 (Pt 9) (1998) 1217–1225.
- [163] T. Hofer, L. Venance, C. Giaume, Control and plasticity of intercellular calcium waves in astrocytes: a modeling approach, *J. Neurosci.* 22 (2002) 4850–4859.
- [164] S. Boitano, E.R. Dirksen, M.J. Sanderson, Intercellular propagation of calcium waves mediated by inositol trisphosphate, *Science* 258 (1992) 292–295.
- [165] H. Niessen, H. Harz, P. Bedner, K. Kramer, K. Willecke, Selective permeability of different connexin channels to the second messenger inositol 1,4,5-trisphosphate, *J. Cell Sci.* 113 (Pt 8) (2000) 1365–1372.
- [166] L. Leybaert, M.J. Sanderson, Intercellular calcium signaling and flash photolysis of caged compounds. A sensitive method to evaluate gap junctional coupling, *Methods Mol. Biol.* 154 (2001) 407–430.
- [167] H. Higashida, A.B. Salmina, R.Y. Olovyanikova, M. Hashii, S. Yokoyama, K. Koizumi, D. Jin, H.X. Liu, O. Lopatina, S. Amina, M.S. Islam, J.J. Huang, M. Noda, Cyclic ADP-ribose as a universal calcium signal molecule in the nervous system, *Neurochem. Int.* 51 (2007) 192–199.
- [168] I.M. Ernst, R. Fliegert, A.H. Guse, Adenine dinucleotide second messengers and T-lymphocyte calcium signaling, *Front. Immunol.* 4 (2013) 259.
- [169] G. Arcuino, J.H. Lin, T. Takano, C. Liu, L. Jiang, Q. Gao, J. Kang, M. Nedergaard, Intercellular calcium signaling mediated by point-source burst release of ATP, *Proc. Natl. Acad. Sci. U. S. A.* 99 (2002) 9840–9845.
- [170] P.B. Guthrie, J. Knappenberger, M. Segal, M.V. Bennett, A.C. Charles, S.B. Kater, ATP released from astrocytes mediates glial calcium waves, *J. Neurosci.* 19 (1999) 520–528.
- [171] C.E. Stout, J.L. Costantin, C.C. Naus, A.C. Charles, Intercellular calcium signaling in astrocytes via ATP release through connexin hemichannels, *J. Biol. Chem.* 277 (2002) 10482–10488.
- [172] N. Hamilton, S. Vayro, F. Kirchhoff, A. Verkhratsky, J. Robbins, D.C. Gorecki, A.M. Butt, Mechanisms of ATP- and glutamate-mediated calcium signaling in white matter astrocytes, *Glia* 56 (2008) 734–749.
- [173] B. Innocenti, V. Pappas, P.G. Haydon, Imaging extracellular waves of glutamate during calcium signaling in cultured astrocytes, *J. Neurosci.* 20 (2000) 1800–1808.
- [174] N.J. Willmott, K. Wong, A.J. Strong, Intercellular  $\text{Ca}^{2+}$  waves in rat hippocampal slice and dissociated glial-neuron cultures mediated by nitric oxide, *FEBS Lett.* 487 (2000) 239–247.
- [175] C. Verderio, S. Bruzzone, E. Zocchi, E. Fedele, U. Schenk, A. De Flora, M. Matteoli, Evidence of a role for cyclic ADP-ribose in calcium signalling and neurotransmitter release in cultured astrocytes, *J. Neurochem.* 78 (2001) 646–657.
- [176] D.N. Bowser, B.S. Khakh, Vesicular ATP is the predominant cause of intercellular calcium waves in astrocytes, *J. Gen. Physiol.* 129 (2007) 485–491.
- [177] N.B. Hamilton, D. Attwell, Do astrocytes really exocytose neurotransmitters? *Nat. Rev. Neurosci.* 11 (2010) 227–238.
- [178] V. Pappas, B.J. Baker, M. Jeras, R. Zorec, Regulated exocytosis in astrocytic signal integration, *Neurochem. Int.* 57 (2010) 451–459.
- [179] M.L. Cotrina, J.H. Lin, A. Alves-Rodrigues, S. Liu, J. Li, H. Azmi-Ghadimi, J. Kang, C.C. Naus, M. Nedergaard, Connexins regulate calcium signaling by controlling ATP release, *Proc. Natl. Acad. Sci. U. S. A.* 95 (1998) 15735–15740.
- [180] A. Baroja-Mazo, M. Barbera-Cremades, P. Pelegrin, The participation of plasma membrane hemichannels to purinergic signaling, *Biochim. Biophys. Acta* 1828 (2013) 79–93.
- [181] S. Locovei, J. Wang, G. Dahl, Activation of pannexin 1 channels by ATP through P2Y receptors and by cytoplasmic calcium, *FEBS Lett.* 580 (2006) 239–244.
- [182] P. Bezzi, V. Gundersen, J.L. Galbete, G. Seifert, C. Steinhäuser, E. Pilati, A. Volterra, Astrocytes contain a vesicular compartment that is competent for regulated exocytosis of glutamate, *Nat. Neurosci.* 7 (2004) 613–620.
- [183] V. Montana, E.B. Malarkey, C. Verderio, M. Matteoli, V. Pappas, Vesicular transmitter release from astrocytes, *Glia* 54 (2006) 700–715.
- [184] A.H. Cornell-Bell, S.M. Finkbeiner, M.S. Cooper, S.J. Smith, Glutamate induces calcium waves in cultured astrocytes: long-range glial signaling, *Science* 247 (1990) 470–473.
- [185] S.J. Smith, Do astrocytes process neural information? *Prog. Brain Res.* 94 (1992) 119–136.
- [186] A.C. Charles, J.E. Merrill, E.R. Dirksen, M.J. Sanderson, Intercellular signaling in glial cells: calcium waves and oscillations in response to mechanical stimulation and glutamate, *Neuron* 6 (1991) 983–992.
- [187] S. Zanotti, A. Charles, Extracellular calcium sensing by glial cells: low extracellular calcium induces intracellular calcium release and intercellular signaling, *J. Neurochem.* 69 (1997) 594–602.
- [188] T.A. Weissman, P.A. Riquelme, L. Ivic, A.C. Flint, A.R. Kriegstein, Calcium waves propagate through radial glial cells and modulate proliferation in the developing neocortex, *Neuron* 43 (2004) 647–661.
- [189] M.E. Harris-White, S.A. Zanotti, S.A. Frautschi, A.C. Charles, Spiral intercellular calcium waves in hippocampal slice cultures, *J. Neurophysiol.* 79 (1998) 1045–1052.
- [190] O. Peters, C.G. Schipke, Y. Hashimoto, H. Kettenmann, Different mechanisms promote astrocyte  $\text{Ca}^{2+}$  waves and spreading depression in the mouse neocortex, *J. Neurosci.* 23 (2003) 9888–9896.
- [191] C.G. Schipke, B. Haas, H. Kettenmann, Astrocytes discriminate and selectively respond to the activity of a subpopulation of neurons within the barrel cortex, *Cereb. Cortex* 18 (2008) 2450–2459.
- [192] X. Wang, T. Takano, M. Nedergaard, Astrocytic calcium signaling: mechanism and implications for functional brain imaging, *Methods Mol. Biol.* 489 (2009) 93–109.
- [193] H. Hirase, L. Qian, P. Bartho, G. Buzsaki, Calcium dynamics of cortical astrocytic networks in vivo, *PLoS Biol.* 2 (2004) E96.
- [194] J. Schummers, H. Yu, M. Sur, Tuned responses of astrocytes and their influence on hemodynamic signals in the visual cortex, *Science* 320 (2008) 1638–1643.
- [195] X. Wang, N. Lou, Q. Xu, G.F. Tian, W.G. Peng, X. Han, J. Kang, T. Takano, M. Nedergaard, Astrocytic  $\text{Ca}^{2+}$  signaling evoked by sensory stimulation in vivo, *Nat. Neurosci.* 9 (2006) 816–823.
- [196] Z.L. Kurth-Nelson, A. Mishra, E.A. Newman, Spontaneous glial calcium waves in the retina develop over early adulthood, *J. Neurosci.* 29 (2009) 11339–11346.
- [197] G.F. Tian, T. Takano, J.H. Lin, X. Wang, L. Bekar, M. Nedergaard, Imaging of cortical astrocytes using 2-photon laser scanning microscopy in the intact mouse brain, *Adv. Drug Deliv. Rev.* 58 (2006) 773–787.
- [198] B. Haas, C.G. Schipke, O. Peters, G. Sohl, K. Willecke, H. Kettenmann, Activity-dependent ATP-waves in the mouse neocortex are independent from astrocytic calcium waves, *Cereb. Cortex* 16 (2006) 237–246.
- [199] E.A. Newman, Propagation of intercellular calcium waves in retinal astrocytes and Muller cells, *J. Neurosci.* 21 (2001) 2215–2223.
- [200] C.G. Schipke, C. Boucsein, C. Ohlemeyer, F. Kirchhoff, H. Kettenmann, Astrocyte  $\text{Ca}^{2+}$  waves trigger responses in microglial cells in brain slices, *FASEB J.* 16 (2002) 255–257.
- [201] A. Araque, R.P. Sanzgiri, V. Pappas, P.G. Haydon, Calcium elevation in astrocytes causes an NMDA receptor-dependent increase in the frequency of miniature synaptic currents in cultured hippocampal neurons, *J. Neurosci.* 18 (1998) 6822–6829.
- [202] E.A. Newman, K.R. Zahs, Modulation of neuronal activity by glial cells in the retina, *J. Neurosci.* 18 (1998) 4022–4028.
- [203] J. Kang, L. Jiang, S.A. Goldman, M. Nedergaard, Astrocyte-mediated potentiation of inhibitory synaptic transmission, *Nat. Neurosci.* 1 (1998) 683–692.
- [204] M. Nedergaard, B. Ransom, S.A. Goldman, New roles for astrocytes: redefining the functional architecture of the brain, *Trends Neurosci.* 26 (2003) 523–530.
- [205] W. Sun, E. McConnell, J.F. Pare, Q. Xu, M. Chen, W. Peng, D. Lovatt, X. Han, Y. Smith, M. Nedergaard, Glutamate-dependent neuroglial calcium signaling differs between young and adult brain, *Science* 339 (2013) 197–200.
- [206] P. Bezzi, G. Carmignoto, L. Pasti, S. Vesce, D. Rossi, B.L. Rizzini, T. Pozzan, A. Volterra, Prostaglandins stimulate calcium-dependent glutamate release in astrocytes, *Nature* 391 (1998) 281–285.
- [207] T. Fellin, O. Pascual, S. Gobbo, T. Pozzan, P.G. Haydon, G. Carmignoto, Neuronal synchrony mediated by astrocytic glutamate through activation of extrasynaptic NMDA receptors, *Neuron* 43 (2004) 729–743.
- [208] M.C. Angulo, A.S. Kozlov, S. Chappak, E. Audinat, Glutamate released from glial cells synchronizes neuronal activity in the hippocampus, *J. Neurosci.* 24 (2004) 6920–6927.
- [209] P.G. Haydon, G. Carmignoto, Astrocyte control of synaptic transmission and neurovascular coupling, *Physiol. Rev.* 86 (2006) 1009–1031.
- [210] G. Dallerac, O. Chever, N. Rouach, How do astrocytes shape synaptic transmission? Insights from electrophysiology, *Front. Cell. Neurosci.* 7 (2013) 159.
- [211] M.A. Di Castro, J. Chuquet, N. Liaudet, K. Bhaukaurally, M. Santello, D. Bouvier, P. Tiret, A. Volterra, Local  $\text{Ca}^{2+}$  detection and modulation of synaptic release by astrocytes, *Nat. Neurosci.* 14 (2011) 1276–1284.
- [212] G.R. Gordon, K.J. Iremonger, S. Kantevari, G.C. Ellis-Davies, B.A. MacVicar, J.S. Bains, Astrocyte-mediated distributed plasticity at hypothalamic glutamate synapses, *Neuron* 64 (2009) 391–403.
- [213] X. Hua, E.B. Malarkey, V. Sunjara, S.E. Rosenwald, W.H. Li, V. Pappas,  $\text{Ca}^{2+}$ -dependent glutamate release involves two classes of endoplasmic reticulum  $\text{Ca}^{2+}$  stores in astrocytes, *J. Neurosci.* 24 (2004) 86–97.
- [214] E.B. Malarkey, Y. Ni, V. Pappas,  $\text{Ca}^{2+}$  entry through TRPC1 channels contributes to intracellular  $\text{Ca}^{2+}$  dynamics and consequent glutamate release from rat astrocytes, *Glia* 56 (2008) 821–835.
- [215] R.C. Reyes, V. Pappas, Mitochondria modulate  $\text{Ca}^{2+}$ -dependent glutamate release from rat cortical astrocytes, *J. Neurosci.* 28 (2008) 9682–9691.
- [216] J. Petrávic, T.A. Fiacco, K.D. McCarthy, Loss of  $\text{IP}_3$  receptor-dependent  $\text{Ca}^{2+}$  increases in hippocampal astrocytes does not affect baseline CA1 pyramidal neuron synaptic activity, *J. Neurosci.* 28 (2008) 4967–4973.

- [217] L.A. Holtzclaw, S. Pandhit, D.J. Bare, G.A. Mignery, J.T. Russell, Astrocytes in adult rat brain express type 2 inositol 1,4,5-trisphosphate receptors, *Glia* 39 (2002) 69–84.
- [218] Q.P. Dong, J.Q. He, Z. Chai, Astrocytic  $\text{Ca}^{2+}$  waves mediate activation of extrasynaptic NMDA receptors in hippocampal neurons to aggravate brain damage during ischemia, *Neurobiol. Dis.* 58 (2013) 68–75.
- [219] S.J. Mulligan, B.A. MacVicar, Calcium transients in astrocyte endfeet cause cerebrovascular constrictions, *Nature* 431 (2004) 195–199.
- [220] K. Paemeleire, L. Leybaert, ATP-dependent astrocyte-endothelial calcium signaling following mechanical damage to a single astrocyte in astrocyte-endothelial cocultures, *J. Neurotrauma* 17 (2000) 345–358.
- [221] G. Burnstock, Purinergic signaling and vascular cell proliferation and death, *Arterioscler. Thromb. Vasc. Biol.* 22 (2002) 364–373.
- [222] S.V. Straub, M.T. Nelson, Astrocytic calcium signaling: the information currency coupling neuronal activity to the cerebral microcirculation, *Trends Cardiovasc. Med.* 17 (2007) 183–190.
- [223] M. Zonta, M.C. Angulo, S. Gobbo, B. Rosengarten, K.A. Hossmann, T. Pozzan, G. Carmignoto, Neuron-to-astrocyte signaling is central to the dynamic control of brain microcirculation, *Nat. Neurosci.* 6 (2003) 43–50.
- [224] T. Takano, G.F. Tian, W. Peng, N. Lou, W. Libionka, X. Han, M. Nedergaard, Astrocyte-mediated control of cerebral blood flow, *Nat. Neurosci.* 9 (2006) 260–267.
- [225] K. Nizar, H. Uhlirava, P. Tian, P.A. Saisan, Q. Cheng, L. Reznichenko, K.L. Weldy, T.C. Steed, V.B. Sridhar, C.L. MacDonald, J. Cui, S.L. Gratiy, S. Sakadzic, D.A. Boas, T.I. Beka, G.T. Einevoll, J. Chen, E. Masliah, A.M. Dale, G.A. Silva, A. Devor, In vivo stimulus-induced vasodilation occurs without IP3 receptor activation and may precede astrocytic calcium increase, *J. Neurosci.* 33 (2013) 8411–8422.
- [226] G.C. Petzold, D.F. Albeanu, T.F. Sato, V.N. Murthy, Coupling of neural activity to blood flow in olfactory glomeruli is mediated by astrocytic pathways, *Neuron* 58 (2008) 897–910.
- [227] R.N. Nishimura, D. Santos, S.T. Fu, B.E. Dwyer, Induction of cell death by L-alphaminoacidic acid exposure in cultured rat astrocytes: relationship to protein synthesis, *Neurotoxicology* 21 (2000) 313–320.
- [228] C.W. Leffler, H. Parfenova, A.L. Fedinec, S. Basuroy, D. Tcheranova, Contributions of astrocytes and CO to pial arteriolar dilation to glutamate in newborn pigs, *Am. J. Physiol. Heart Circ. Physiol.* 291 (2006) H2897–H2904.
- [229] Q. Xi, E. Umstot, G. Zhao, D. Narayanan, C.W. Leffler, J.H. Jaggar, Glutamate regulates  $\text{Ca}^{2+}$  signals in smooth muscle cells of newborn piglet brain slice arterioles through astrocyte- and heme oxygenase-dependent mechanisms, *Am. J. Physiol. Heart Circ. Physiol.* 298 (2010) H562–H569.
- [230] D.L. Price, J.W. Ludwig, H. Mi, T.L. Schwarz, M.H. Ellisman, Distribution of rSlo  $\text{Ca}^{2+}$ -activated  $\text{K}^{+}$  channels in rat astrocyte perivascular endfeet, *Brain Res.* 956 (2002) 183–193.
- [231] H. Girouard, A.D. Bonev, R.M. Hannah, A. Meredith, R.W. Aldrich, M.T. Nelson, Astrocytic endfoot  $\text{Ca}^{2+}$  and BK channels determine both arteriolar dilation and constriction, *Proc. Natl. Acad. Sci. U. S. A.* 107 (2010) 3811–3816.
- [232] M. Koide, A.D. Bonev, M.T. Nelson, G.C. Wellman, Inversion of neurovascular coupling by subarachnoid blood depends on large-conductance  $\text{Ca}^{2+}$ -activated  $\text{K}^{+}$  (BK) channels, *Proc. Natl. Acad. Sci. U. S. A.* 109 (2012) E1387–E1395.
- [233] R.G. Gordon, H.B. Choi, R.L. Runge, G.C. Ellis-Davies, B.A. MacVicar, Brain metabolism dictates the polarity of astrocyte control over arterioles, *Nature* 456 (2008) 745–749.
- [234] K.M. Dunn, D.C. Hill-Eubanks, W.B. Liedtke, M.T. Nelson, TRPV4 channels stimulate  $\text{Ca}^{2+}$ -induced  $\text{Ca}^{2+}$  release in astrocytic endfeet and amplify neurovascular coupling responses, *Proc. Natl. Acad. Sci. U. S. A.* 110 (2013) 6157–6162.
- [235] H. Girouard, C. Iadecola, Neurovascular coupling in the normal brain and in hypertension, stroke, and Alzheimer disease, *J. Appl. Physiol.* 100 (2006) 328–335.
- [236] M. Belanger, I. Allaman, P.J. Magistretti, Brain energy metabolism: focus on astrocyte–neuron metabolic cooperation, *Cell Metab.* 14 (2011) 724–738.
- [237] K.A. Kasischke, H.D. Vishwasrao, P.J. Fisher, W.R. Zipfel, W.W. Webb, Neural activity triggers neuronal oxidative metabolism followed by astrocytic glycolysis, *Science* 305 (2004) 99–103.
- [238] L. Pellerin, A.K. Bouzier-Sore, A. Aubert, S. Serres, M. Merle, R. Costalat, P.J. Magistretti, Activity-dependent regulation of energy metabolism by astrocytes: an update, *Glia* 55 (2007) 1251–1262.
- [239] L. Pellerin, G. Pellegrini, P.G. Bittar, Y. Charnay, C. Bouras, J.L. Martin, N. Stella, P.J. Magistretti, Evidence supporting the existence of an activity-dependent astrocyte–neuron lactate shuttle, *Dev. Neurosci.* 20 (1998) 291–299.
- [240] L. Leybaert, Neurobarrier coupling in the brain: a partner of neurovascular and neurometabolic coupling? *J. Cereb. Blood Flow Metab.* 25 (2005) 2–16.
- [241] U. Rabast, E. Vogelhuber, K.H. Vornberger, M. Ehl, H. Kasper, Long-term studies in obese patients after hospital and ambulatory carbohydrate-reducing diet therapy, *Verh. Dtsch. Ges. Inn. Med.* (1978) 1151–1154.
- [242] R.K. Prasad, F. Ismail-Beigi, Mechanism of stimulation of glucose transport by H2O2: role of phospholipase C, *Arch. Biochem. Biophys.* 362 (1999) 113–122.
- [243] R.A. Quintanilla, O.H. Porras, J. Castro, L.F. Barros, Cytosolic  $\text{Ca}^{2+}$  modulates basal GLUT1 activity and plays a permissive role in its activation by metabolic stress and insulin in rat epithelial cells, *Cell Calcium* 28 (2000) 97–106.
- [244] P. Singh, J.J. Leddy, G.J. Chatzys, M. Salih, B.S. Tuana, Alternative splicing generates a CaM kinase II $\beta$  isoform in myocardium that targets the sarcoplasmic reticulum through a putative  $\alpha$ KAP and regulates GAPDH, *Mol. Cell. Biochem.* 270 (2005) 215–221.
- [245] T. Pancani, K.L. Anderson, N.M. Porter, O. Thibault, Imaging of a glucose analog, calcium and NADH in neurons and astrocytes: dynamic responses to depolarization and sensitivity to pioglitazone, *Cell Calcium* 50 (2011) 548–558.
- [246] T.A. Fiacco, C. Agulhon, K.D. McCarthy, Sorting out astrocyte physiology from pharmacology, *Annu. Rev. Pharmacol. Toxicol.* 49 (2009) 151–174.
- [247] J. Chuquet, L. Hollender, E.A. Nimchinsky, High-resolution in vivo imaging of the neurovascular unit during spreading depression, *J. Neurosci.* 27 (2007) 4036–4044.
- [248] C. Stout, A. Charles, Modulation of intercellular calcium signaling in astrocytes by extracellular calcium and magnesium, *Glia* 43 (2003) 265–273.
- [249] T. Kristian, G. Gido, B.K. Siesjö, Brain calcium metabolism in hypoglycemic coma, *J. Cereb. Blood Flow Metab.* 13 (1993) 955–961.
- [250] A. Vezzani, H.Q. Wu, P. Angelico, M.A. Stasi, R. Samanin, Quinolinic acid-induced seizures, but not nerve cell death, are associated with extracellular  $\text{Ca}^{2+}$  decrease assessed in the hippocampus by brain dialysis, *Brain Res. Brain Res. Rev.* 454 (1988) 289–297.
- [251] G.G. Somjen, P.G. Aitken, J.L. Giacchino, J.O. McNamara, Interstitial ion concentrations and paroxysmal discharges in hippocampal formation and spinal cord, *Adv. Neurol.* 44 (1986) 663–680.
- [252] L. Leybaert, G. De Ley, Interstitial and tissue cations and electrical potential after experimental spinal cord injury, *Exp. Brain Res.* 100 (1994) 369–375.
- [253] A.M. Hofer, Another dimension to calcium signaling: a look at extracellular calcium, *J. Cell Sci.* 118 (2005) 855–862.
- [254] A. Torres, F. Wang, Q. Xu, T. Fujita, R. Dobrowolski, K. Willecke, T. Takano, M. Nedergaard, Extracellular  $\text{Ca}^{2+}$  acts as a mediator of communication from neurons to glia, *Sci. Signal.* 5 (2012) ra8.
- [255] L.R. Mills, A.A. Velumian, S.K. Agrawal, E. Theriault, M.G. Fehlings, Confocal imaging of changes in glial calcium dynamics and homeostasis after mechanical injury in rat spinal cord white matter, *Neuroimage* 21 (2004) 1069–1082.
- [256] S.E. Samuels, J.B. Lipitz, G. Dahl, K.J. Muller, Neuroglial ATP release through innexin channels controls microglial cell movement to a nerve injury, *J. Gen. Physiol.* 136 (2010) 425–442.
- [257] A. Nimmerjahn, E.A. Mukamel, M.J. Schnitzer, Motor behavior activates Bergmann glial networks, *Neuron* 62 (2009) 400–412.
- [258] N.J. Haughey, M.P. Mattson, Alzheimer's amyloid  $\beta$ -peptide enhances ATP/gap junction-mediated calcium-wave propagation in astrocytes, *Neuromol. Med.* 3 (2003) 173–180.
- [259] S.K. Chow, D. Yu, C.L. Macdonald, M. Buibas, G.A. Silva, Amyloid  $\beta$ -peptide directly induces spontaneous calcium transients, delayed intercellular calcium waves and gliosis in rat cortical astrocytes, *ASN Neuro* 2 (2010) e00026.
- [260] L. Lee, P. Kosuri, O. Arancio, Picomolar amyloid- $\beta$  peptides enhance spontaneous astrocyte calcium transients, *J. Alzheimers Dis.* 38 (2014) 49–62.
- [261] M.A. Busche, G. Eichhoff, H. Adelsberger, D. Abramowski, K.H. Wiederhold, C. Haass, M. Staufenbiel, A. Konnerth, O. Garaschuk, Clusters of hyperactive neurons near amyloid plaques in a mouse model of Alzheimer's disease, *Science* 321 (2008) 1686–1689.
- [262] E. Alberdi, A. Wyssenbach, M. Alberdi, M.V. Sanchez-Gomez, F. Cavaliere, J.J. Rodriguez, A. Verkhratsky, C. Matute,  $\text{Ca}^{2+}$ -dependent endoplasmic reticulum stress correlates with astrogliosis in oligomeric amyloid  $\beta$ -treated astrocytes and in a model of Alzheimer's disease, *Aging Cell* 12 (2013) 292–302.
- [263] U. Armato, A. Chiarini, B. Chakravarthy, F. Chioffi, R. Pacchiana, E. Colarusso, J.F. Whitfield, I. Dal Pra, Calcium-sensing receptor antagonist (calcilytic) NPS 2143 specifically blocks the increased secretion of endogenous Abeta42 prompted by exogenous fibrillary or soluble Abeta25–35 in human cortical astrocytes and neurons—therapeutic relevance to Alzheimer's disease, *Biochim. Biophys. Acta* 1832 (2013) 1634–1652.
- [264] M.H. Sarker, D.E. Hu, P.A. Fraser, Acute effects of bradykinin on cerebral microvascular permeability in the anesthetized rat, *J. Physiol.* 528 (Pt 1) (2000) 177–187.
- [265] M. Kawahara, I. Ohtsuka, S. Yokoyama, M. Kato-Negishi, Y. Sadakane, Membrane incorporation, channel formation, and disruption of calcium homeostasis by Alzheimer's  $\beta$ -amyloid protein, *Int. J. Alzheimers Dis.* 2011 (2011) 304583.
- [266] G. Burnstock, U. Krugel, M.P. Abbracchio, P. Illes, Purinergic signalling: from normal behaviour to pathological brain function, *Prog. Neurobiol.* 95 (2011) 229–274.
- [267] E. Decrock, E. De Vuyst, M. Vinken, M. Van Moorhem, K. Vranckx, N. Wang, L. Van Laeken, M. De Bock, K. D'Herde, C.P. Lai, V. Rogiers, W.H. Evans, C.C. Naus, L. Leybaert, Connexin 43 hemichannels contribute to the propagation of apoptotic cell death in a rat C6 glioma cell model, *Cell Death Differ.* 16 (2009) 151–163.
- [268] E. Decrock, D.V. Krysko, M. Vinken, A. Kaczmarek, G. Crispino, M. Bol, N. Wang, M. De Bock, E. De Vuyst, C.C. Naus, V. Rogiers, P. Vandenabeele, C. Erneux, F. Mammano, G. Bultynck, L. Leybaert, Transfer of IP(3) through gap junctions is critical, but not sufficient, for the spread of apoptosis, *Cell Death Differ.* 19 (2012) 947–957.
- [269] E. Decrock, M. De Bock, N. Wang, A.K. Gadicherla, M. Bol, T. Delvaeye, P. Vandenabeele, M. Vinken, G. Bultynck, D.V. Krysko, L. Leybaert, IP3, a small molecule with a powerful message, *Biochim. Biophys. Acta* 1833 (2013) 1772–1786.
- [270] E.A. Eugenini, J.W. Berman, Cytochrome c dysregulation induced by HIV infection of astrocytes results in bystander apoptosis of uninfected astrocytes by an IP and calcium-dependent mechanism, *J. Neurochem.* 127 (2013) 644–651.
- [271] J.P. Decuyper, G. Monaco, G. Bultynck, L. Missiaen, H. De Smedt, J.B. Parys, The IP(3) receptor–mitochondria connection in apoptosis and autophagy, *Biochim. Biophys. Acta* 1813 (2011) 1003–1013.
- [272] C.C. Mendes, D.A. Gomes, M. Thompson, N.C. Souto, T.S. Goes, A.M. Goes, M.A. Rodrigues, M.V. Gomez, M.H. Nathanson, M.F. Leite, The type III inositol 1,4,5-trisphosphate receptor preferentially transmits apoptotic  $\text{Ca}^{2+}$  signals into mitochondria, *J. Biol. Chem.* 280 (2005) 40892–40900.
- [273] D. De Stefani, A. Bononi, A. Romagnoli, A. Messina, V. De Pinto, P. Pinton, R. Rizzuto, VDAC1 selectively transfers apoptotic  $\text{Ca}^{2+}$  signals to mitochondria, *Cell Death Differ.* 19 (2012) 267–273.
- [274] P. Pinton, C. Giorgi, R. Síviero, E. Zecchini, R. Rizzuto, Calcium and apoptosis: ER–mitochondria  $\text{Ca}^{2+}$  transfer in the control of apoptosis, *Oncogene* 27 (2008) 6407–6418.

- [275] C. Giorgi, M.R. Wieckowski, P.P. Pandolfi, P. Pinton, Mitochondria associated membranes (MAMs) as critical hubs for apoptosis, *Commun. Integr. Biol.* 4 (2011) 334–335.
- [276] S. Grimm, The ER-mitochondria interface: the social network of cell death, *Biochim. Biophys. Acta* 1823 (2012) 327–334.
- [277] S. Kiviluoto, H. Akl, T. Vervliet, G. Bultynck, J.B. Parys, L. Missiaen, H. De Smedt, IP3 receptor-binding partners in cell-death mechanisms, *Wiley Interdiscip. Rev. Membr. Transp. Signal* 1 (2012) 201–210.
- [278] G. Monaco, T. Vervliet, H. Akl, G. Bultynck, The selective BH4-domain biology of Bcl-2-family members: IP3Rs and beyond, *Cell. Mol. Life Sci.* 70 (2013) 1171–1183.
- [279] Y.B. Ouyang, R.G. Giffard, ER-mitochondria crosstalk during cerebral ischemia: molecular chaperones and ER-mitochondrial calcium transfer, *Int. J. Cell Biol.* 2012 (2012) 493934.
- [280] Y.B. Ouyang, L.J. Xu, J.F. Emery, A.S. Lee, R.G. Giffard, Overexpressing GRP78 influences Ca<sup>2+</sup> handling and function of mitochondria in astrocytes after ischemia-like stress, *Mitochondrion* 11 (2011) 279–286.
- [281] L.E. Jensen, G. Bultynck, T. Luyten, H. Amijee, M.D. Bootman, H.L. Roderick, Alzheimer's disease-associated peptide Abeta42 mobilizes ER Ca<sup>2+</sup> via InsP3R-dependent and -independent mechanisms, *Front. Mol. Neurosci.* 6 (2013) 36.
- [282] M.J. Kipanyula, L. Contreras, E. Zampese, C. Lazzari, A.K. Wong, P. Pizzo, C. Fasolato, T. Pozzan, Ca<sup>2+</sup> dysregulation in neurons from transgenic mice expressing mutant presenilin 2, *Aging Cell* 11 (2013) 885–893.
- [283] K.H. Cheung, D. Shineman, M. Muller, C. Cardenas, L. Mei, J. Yang, T. Tomita, T. Iwatsubo, V.M. Lee, J.K. Foskett, Mechanism of Ca<sup>2+</sup> disruption in Alzheimer's disease by presenilin regulation of InsP3 receptor channel gating, *Neuron* 58 (2008) 871–883.
- [284] T. Higo, K. Hamada, C. Hisatsune, N. Nukina, T. Hashikawa, M. Hattori, T. Nakamura, K. Mikoshiba, Mechanism of ER stress-induced brain damage by IP(3) receptor, *Neuron* 68 (2010) 865–878.
- [285] T. Cali, D. Ottolini, A. Negro, M. Brini, Enhanced parkin levels favor ER-mitochondria crosstalk and guarantee Ca<sup>2+</sup> transfer to sustain cell bioenergetics, *Biochim. Biophys. Acta* 1832 (2013) 495–508.
- [286] T. Cali, D. Ottolini, A. Negro, M. Brini, alpha-Synuclein controls mitochondrial calcium homeostasis by enhancing endoplasmic reticulum-mitochondria interactions, *J. Biol. Chem.* 287 (2012) 17914–17929.
- [287] W. Zheng, L. Talley Watts, D.M. Holstein, J. Wewer, J.D. Lechleiter, P2Y1R-initiated, IP3R-dependent stimulation of astrocyte mitochondrial metabolism reduces and partially reverses ischemic neuronal damage in mouse, *J. Cereb. Blood Flow Metab.* 33 (2013) 600–611.
- [288] G. Begum, D. Kintner, Y. Liu, S.W. Cramer, D. Sun, DHA inhibits ER Ca<sup>2+</sup> release and ER stress in astrocytes following in vitro ischemia, *J. Neurochem.* 120 (2012) 622–630.
- [289] J.M. Johnston, P. Burnett, A.P. Thomas, N. Tezapsidis, Calcium oscillations in type-1 astrocytes, the effect of a presenilin 1 (PS1) mutation, *Neurosci. Lett.* 395 (2006) 159–164.
- [290] A.A. Grolla, J.A. Sim, D. Lim, J.J. Rodriguez, A.A. Genazzani, A. Verkhratsky, Amyloid-beta and Alzheimer's disease type pathology differentially affects the calcium signalling toolkit in astrocytes from different brain regions, *Cell Death Dis.* 4 (2013) e623.
- [291] E. Decrock, M. Vinken, E. De Vuyst, D.V. Krysko, K. D'Herde, T. Vanhaecke, P. Vandenabeele, V. Rogiers, L. Leybaert, Connexin-related signaling in cell death: to live or let die? *Cell Death Differ.* 16 (2009) 524–536.
- [292] E. Decrock, M. Vinken, M. Bol, K. D'Herde, V. Rogiers, P. Vandenabeele, D.V. Krysko, G. Bultynck, L. Leybaert, Calcium and connexin-based intercellular communication, a deadly catch? *Cell Calcium* 50 (2011) 310–321.
- [293] M.E. Hamby, M.V. Sofroniew, Reactive astrocytes as therapeutic targets for CNS disorders, *Neurotherapeutics* 7 (2010) 494–506.
- [294] I. Fernaldo-Espinosa, M. Nieto-Sampedro, P. Bovolenta, Differential activation of microglia and astrocytes in aniso- and isomorphic gliotic tissue, *Glia* 8 (1993) 277–291.
- [295] J. Silver, J.H. Miller, Regeneration beyond the glial scar, *Nat. Rev. Neurosci.* 5 (2004) 146–156.
- [296] M. Pekny, M. Nilsson, Astrocyte activation and reactive gliosis, *Glia* 50 (2005) 427–434.
- [297] M.V. Sofroniew, Molecular dissection of reactive astrogliosis and glial scar formation, *Trends Neurosci.* 32 (2009) 638–647.
- [298] M.E. Hamby, G. Coppola, Y. Ao, D.H. Geschwind, B.S. Khakh, M.V. Sofroniew, Inflammatory mediators alter the astrocyte transcriptome and calcium signaling elicited by multiple G-protein-coupled receptors, *J. Neurosci.* 32 (2012) 14489–14510.
- [299] O. Butenko, D. Dzamba, J. Benesova, P. Honsa, V. Benfenati, V. Rusnakova, S. Ferroni, M. Anderova, The increased activity of TRPV4 channel in the astrocytes of the adult rat hippocampus after cerebral hypoxia/ischemia, *PLoS One* 7 (2012) e39959.
- [300] N. Daschil, G.J. Obermair, B.E. Flucher, N. Stefanova, B. Hutter-Paier, M. Windisch, C. Humpel, J. Marksteiner, Cav1.2 calcium channel expression in reactive astrocytes is associated with the formation of amyloid-beta plaques in an Alzheimer's disease mouse model, *J. Alzheimers Dis.* 37 (2013) 439–451.
- [301] J.H. Li, S.T. Zhao, C.Y. Wu, X. Cao, M.R. Peng, S.J. Li, X.A. Liu, T.M. Gao, Store-operated Ca<sup>2+</sup> channels blockers inhibit lipopolysaccharide induced astrocyte activation, *Neurochem. Res.* 38 (2013) 2216–2226.
- [302] K. Kanamaru, J. Kubota, H. Sekiya, K. Hirose, Y. Okubo, M. Iino, Calcium-dependent N-cadherin up-regulation mediates reactive astrogliosis and neuroprotection after brain injury, *Proc. Natl. Acad. Sci. U. S. A.* 110 (2013) 11612–11617.
- [303] S. Du, A. Rubin, S. Klepper, C. Barrett, Y.C. Kim, H.W. Rhim, E.B. Lee, C.W. Park, G.J. Markelonis, T.H. Oh, Calcium influx and activation of calpain I mediate acute reactive gliosis in injured spinal cord, *Exp. Neurol.* 157 (1999) 96–105.
- [304] K. Gao, C.R. Wang, F. Jiang, A.Y. Wong, N. Su, J.H. Jiang, R.C. Chai, G. Vatcher, J. Teng, J. Chen, Y.W. Jiang, A.C. Yu, Traumatic scratch injury in astrocytes triggers calcium influx to activate the JNK/c-Jun/AP-1 pathway and switch on GFAP expression, *Glia* 61 (2013) 2063–2077.
- [305] N. Theodoris, J.F. Bechberger, C.C. Naus, W.C. Sin, Role of gap junction protein connexin43 in astrogliosis induced by brain injury, *PLoS One* 7 (2012) e47311.
- [306] Z. Wu, H. Xu, Y. He, G. Yang, C. Liao, W. Gao, M. Liang, X. He, Antisense oligodeoxynucleotides targeting connexin43 reduce cerebral astrocytosis and edema in a rat model of traumatic brain injury, *Neurol. Res.* 35 (2013) 255–262.
- [307] X. Zhang, B. Cao, J. Wang, J. Liu, V.O. Tung, P.K. Lam, L.L. Chan, Y. Li, Neurotoxicity and reactive astrogliosis in the anterior cingulate cortex in acute ciguatera poisoning, *Neuromol. Med.* 15 (2013) 310–323.
- [308] A. Ohsumi, H. Nawashiro, N. Otani, H. Oigawa, T. Toyooka, A. Yano, N. Nomura, K. Shima, Alteration of gap junction proteins (connexins) following lateral fluid percussion injury in rats, *Acta Neurochir. Suppl.* 96 (2006) 148–150.
- [309] M. Cronin, P.N. Anderson, J.E. Cook, C.R. Green, D.L. Becker, Blocking connexin43 expression reduces inflammation and improves functional recovery after spinal cord injury, *Mol. Cell. Neurosci.* 39 (2008) 152–160.
- [310] E. Theriault, U.N. Frankenstein, E.L. Hertzberg, J.I. Nagy, Connexin43 and astrocytic gap junctions in the rat spinal cord after acute compression injury, *J. Comp. Neurol.* 382 (1997) 199–214.
- [311] C. Haupt, O.W. Witte, C. Frahm, Up-regulation of Connexin43 in the glial scar following photothrombotic ischemic injury, *Mol. Cell. Neurosci.* 35 (2007) 89–99.
- [312] M.G. Kozoriz, J.F. Bechberger, G.R. Bechberger, M.W. Suen, A.P. Moreno, K. Maass, K. Willecke, C.C. Naus, The connexin43 C-terminal region mediates neuroprotection during stroke, *J. Neuropathol. Exp. Neurol.* 69 (2010) 196–206.
- [313] X. Mei, P. Ezan, C. Giaume, A. Koulakoff, Astroglial connexin immunoreactivity is specifically altered at beta-amyloid plaques in beta-amyloid precursor protein/presenilin1 mice, *Neuroscience* 171 (2010) 92–105.
- [314] C.G. Fonseca, C.R. Green, L.F. Nicholson, Upregulation in astrocytic connexin 43 gap junction levels may exacerbate generalized seizures in mesial temporal lobe epilepsy, *Brain Res.* 929 (2002) 105–116.
- [315] D.K. Takahashi, J.R. Vargas, K.S. Wilcox, Increased coupling and altered glutamate transport currents in astrocytes following kainic-acid-induced status epilepticus, *Neurobiol. Dis.* 40 (2010) 573–585.
- [316] H.V. Danesh-Meyer, N.M. Kerr, J. Zhang, E.K. Eady, S.J. O'Carroll, L.F. Nicholson, C.S. Johnson, C.R. Green, Connexin43 mimetic peptide reduces vascular leak and retinal ganglion cell death following retinal ischaemia, *Brain* 135 (2012) 506–520.
- [317] N.M. Kerr, C.S. Johnson, J. Zhang, E.K. Eady, C.R. Green, H.V. Danesh-Meyer, High pressure-induced retinal ischaemia reperfusion causes upregulation of gap junction protein connexin43 prior to retinal ganglion cell loss, *Exp. Neurol.* 234 (2012) 144–152.
- [318] C. Huang, X. Han, X. Li, E. Lam, W. Peng, N. Lou, A. Torres, M. Yang, J.M. Garre, G.F. Tian, M.V. Bennett, M. Nedergaard, T. Takano, Critical role of connexin 43 in secondary expansion of traumatic spinal cord injury, *J. Neurosci.* 32 (2012) 3333–3338.
- [319] S.J. O'Carroll, C.A. Gorrie, S. Velamoor, C.R. Green, L.F. Nicholson, Connexin43 mimetic peptide is neuroprotective and improves function following spinal cord injury, *Neurosci. Res.* 75 (2013) 256–267.
- [320] M.J. Chen, B. Kress, X. Han, K. Moll, W. Peng, R.R. Ji, M. Nedergaard, Astrocytic CX43 hemichannels and gap junctions play a crucial role in development of chronic neuropathic pain following spinal cord injury, *Glia* 60 (2012) 1660–1670.
- [321] I.H. Lee, E. Lindqvist, O. Kiehn, J. Widenfalk, L. Olson, Glial and neuronal connexin expression patterns in the rat spinal cord during development and following injury, *J. Comp. Neurol.* 489 (2005) 1–10.
- [322] E. Brand-Schieber, P. Werner, D.A. Iacobas, S. Iacobas, M. Beelitz, S.L. Lowery, D.C. Spray, E. Scemes, Connexin43, the major gap junction protein of astrocytes, is down-regulated in inflamed white matter in an animal model of multiple sclerosis, *J. Neurosci. Res.* 80 (2005) 798–808.
- [323] E. Gangoso, P. Ezan, J.C. Valle-Casuso, S. Herrero-Gonzalez, A. Koulakoff, J.M. Medina, C. Giaume, A. Tabernero, Reduced connexin43 expression correlates with c-Src activation, proliferation, and glucose uptake in reactive astrocytes after an excitotoxic insult, *Glia* 60 (2012) 2040–2049.
- [324] A. Zappala, G. Li Volti, M.F. Serapide, R. Pellitteri, M. Falchi, F. La Delia, V. Cicirata, F. Cicirata, Expression of pannexin2 protein in healthy and ischemized brain of adult rats, *Neuroscience* 148 (2007) 653–667.
- [325] A. Mehta, M. Prabhakar, P. Kumar, R. Deshmukh, P.L. Sharma, Excitotoxicity: bridge to various triggers in neurodegenerative disorders, *Eur. J. Pharmacol.* 698 (2013) 6–18.
- [326] M. Facheris, S. Beretta, C. Ferrarese, Peripheral markers of oxidative stress and excitotoxicity in neurodegenerative disorders: tools for diagnosis and therapy? *J. Alzheimers Dis.* 6 (2004) 177–184.
- [327] N.M. Ashpole, A.R. Chawla, M.P. Martin, T. Brustovetsky, N. Brustovetsky, A. Hudmon, Loss of calcium/calmodulin-dependent protein kinase II activity in cortical astrocytes decreases glutamate uptake and induces neurotoxic release of ATP, *J. Biol. Chem.* 288 (2013) 14599–14611.
- [328] M.V. Frantseva, L. Kokorotseva, J.L. Perez Velazquez, Ischemia-induced brain damage depends on specific gap-junctional coupling, *J. Cereb. Blood Flow Metab.* 22 (2002) 453–462.
- [329] J.L. Perez Velazquez, M.V. Frantseva, C.C. Naus, Gap junctions and neuronal injury: protectants or executioners? *Neuroscientist* 9 (2003) 5–9.
- [330] J.J. O'Connor, Targeting tumour necrosis factor-alpha in hypoxia and synaptic signalling, *Ir. J. Med. Sci.* 182 (2013) 157–162.
- [331] N. Froger, J.A. Orellana, C.F. Calvo, E. Amigou, M.G. Kozoriz, C.C. Naus, J.C. Saez, C. Giaume, Inhibition of cytokine-induced connexin43 hemichannel activity in astrocytes is neuroprotective, *Mol. Cell. Neurosci.* 45 (2010) 37–46.



- [332] E. Salas, L.M. Carrasquero, L.A. Olivos-Ore, D. Bustillo, A.R. Artalejo, M.T. Miras-Portugal, E.G. Delicado, Purinergic P2x7 receptors mediate cell death in mouse cerebellar astrocytes in culture, *J. Pharmacol. Exp. Ther.* 347 (2013) 802–815.
- [333] O. Pascual, K.B. Casper, C. Kubera, J. Zhang, R. Revilla-Sanchez, J.Y. Sul, H. Takano, S.J. Moss, K. McCarthy, P.G. Haydon, Astrocytic purinergic signaling coordinates synaptic networks, *Science* 310 (2005) 113–116.
- [334] M. Domercq, A. Perez-Samartin, D. Aparicio, E. Alberdi, O. Pampliega, C. Matute, P2X7 receptors mediate ischemic damage to oligodendrocytes, *Glia* 58 (2010) 730–740.
- [335] S. Locovei, E. Scemes, F. Qiu, D.C. Spray, G. Dahl, Pannexin1 is part of the pore forming unit of the P2X(7) receptor death complex, *FEBS Lett.* 581 (2007) 483–488.
- [336] P. Pelegrin, A. Surprenant, Pannexin-1 mediates large pore formation and interleukin-1 $\beta$  release by the ATP-gated P2X7 receptor, *EMBO J.* 25 (2006) 5071–5082.
- [337] D. Dzamba, P. Honsa, M. Anderova, NMDA receptors in glial cells: pending questions, *Curr. Neuropharmacol.* 11 (2013) 250–262.
- [338] M. Murugan, E.A. Ling, C. Kaur, Dysregulated glutamate uptake by astrocytes causes oligodendroglia death in hypoxic periventricular white matter damage, *Mol. Cell. Neurosci.* 56 (2013) 342–354.
- [339] C. Farina, F. Aloisi, E. Meinl, Astrocytes are active players in cerebral innate immunity, *Trends Immunol.* 28 (2007) 138–145.
- [340] Y. Dong, E.N. Benveniste, Immune function of astrocytes, *Glia* 36 (2001) 180–190.
- [341] L.T. Lau, A.C. Yu, Astrocytes produce and release interleukin-1, interleukin-6, tumor necrosis factor  $\alpha$  and interferon- $\gamma$  following traumatic and metabolic injury, *J. Neurotrauma* 18 (2001) 351–359.
- [342] C.M. Liberto, P.J. Albrecht, L.M. Herx, V.W. Yong, S.W. Levison, Pro-regenerative properties of cytokine-activated astrocytes, *J. Neurochem.* 89 (2004) 1092–1100.
- [343] T. Ida, M. Hara, Y. Nakamura, S. Kozaki, S. Tsunoda, H. Ihara, Cytokine-induced enhancement of calcium-dependent glutamate release from astrocytes mediated by nitric oxide, *Neurosci. Lett.* 432 (2008) 232–236.
- [344] J.A. Orellana, N. Froger, P. Ezan, J.X. Jiang, M.V. Bennett, C.C. Naus, C. Giaume, J.C. Saez, ATP and glutamate released via astroglial connexin 43 hemichannels mediate neuronal death through activation of pannexin 1 hemichannels, *J. Neurochem.* 118 (2011) 826–840.
- [345] J. Minkiewicz, J.P. de Rivero Vaccari, R.W. Keane, Human astrocytes express a novel NLRP2 inflammasome, *Glia* 61 (2013) 1113–1121.
- [346] X.F. Meng, X.L. Wang, X.J. Tian, Z.H. Yang, G.P. Chu, J. Zhang, M. Li, J. Shi, C. Zhang, Nod-like receptor protein 1 inflammasome mediates neuron injury under high glucose, *Mol. Neurobiol.* 49 (2014) 673–684.
- [347] U.K. Hanisch, H. Kettenmann, Microglia: active sensor and versatile effector cells in the normal and pathologic brain, *Nat. Neurosci.* 10 (2007) 1387–1394.
- [348] V.H. Perry, J.A. Nicoll, C. Holmes, Microglia in neurodegenerative disease, *Nat. Rev. Neurol.* 6 (2010) 193–201.
- [349] H. Kettenmann, U.K. Hanisch, M. Noda, A. Verkhratsky, Physiology of microglia, *Physiol. Rev.* 91 (2011) 461–553.
- [350] R.M. Ransohoff, V.H. Perry, Microglial physiology: unique stimuli, specialized responses, *Annu. Rev. Immunol.* 27 (2009) 119–145.
- [351] M. Noda, M. Ifuku, Y. Mori, A. Verkhratsky, Calcium influx through reversed NCX controls migration of microglia, *Adv. Exp. Med. Biol.* 961 (2013) 289–294.
- [352] K. Farber, H. Kettenmann, Purinergic signaling and microglia, *Pflugers Arch.* 452 (2006) 615–621.
- [353] M. Lee, Neurotransmitters and microglial-mediated neuroinflammation, *Curr. Protein Pept. Sci.* 14 (2013) 21–32.
- [354] K. Farber, H. Kettenmann, Functional role of calcium signals for microglial function, *Glia* 54 (2006) 656–665.
- [355] S.W. Barger, A.S. Basile, Activation of microglia by secreted amyloid precursor protein evokes release of glutamate by cystine exchange and attenuates synaptic function, *J. Neurochem.* 76 (2001) 846–854.
- [356] J.M. Sanz, P. Chiozzi, D. Ferrari, M. Colaianna, M. Idzko, S. Falzoni, R. Fellin, L. Trabace, F. Di Virgilio, Activation of microglia by amyloid ( $\beta$ ) requires P2X7 receptor expression, *J. Immunol.* 182 (2009) 4378–4385.
- [357] T. Mika, N. Prochnow, Functions of connexins and large pore channels on microglial cells: the gates to environment, *Brain Res.* 1487 (2012) 16–24.
- [358] H. Takeuchi, S. Jin, J. Wang, G. Zhang, J. Kawanokuchi, R. Kuno, Y. Sonobe, T. Mizuno, A. Suzumura, Tumor necrosis factor- $\alpha$  induces neurotoxicity via glutamate release from hemichannels of activated microglia in an autocrine manner, *J. Biol. Chem.* 281 (2006) 21362–21368.
- [359] H. Takeuchi, S. Jin, H. Suzuki, Y. Doi, J. Liang, J. Kawanokuchi, T. Mizuno, M. Sawada, A. Suzumura, Blockade of microglial glutamate release protects against ischemic brain injury, *Exp. Neurol.* 214 (2008) 144–146.
- [360] H. Takeuchi, H. Mizoguchi, Y. Doi, S. Jin, M. Noda, J. Liang, H. Li, Y. Zhou, R. Mori, S. Yasuoka, E. Li, B. Parajuli, J. Kawanokuchi, Y. Sonobe, J. Sato, K. Yamanaka, G. Sobue, T. Mizuno, A. Suzumura, Blockade of gap junction hemichannel suppresses disease progression in mouse models of amyotrophic lateral sclerosis and Alzheimer's disease, *PLoS One* 6 (2011) e21108.
- [361] J. Shijie, H. Takeuchi, I. Yawata, Y. Harada, Y. Sonobe, Y. Doi, J. Liang, L. Hua, S. Yasuoka, Y. Zhou, M. Noda, J. Kawanokuchi, T. Mizuno, A. Suzumura, Blockade of glutamate release from microglia attenuates experimental autoimmune encephalomyelitis in mice, *Tohoku J. Exp. Med.* 217 (2009) 87–92.
- [362] J. Liang, H. Takeuchi, Y. Doi, J. Kawanokuchi, Y. Sonobe, S. Jin, I. Yawata, H. Li, S. Yasuoka, T. Mizuno, A. Suzumura, Excitatory amino acid transporter expression by astrocytes is neuroprotective against microglial excitotoxicity, *Brain Res.* 1210 (2008) 11–19.
- [363] J. Takaki, K. Fujimori, M. Miura, T. Suzuki, Y. Sekino, K. Sato, L-glutamate released from activated microglia downregulates astrocytic L-glutamate transporter expression in neuroinflammation: the 'collusion' hypothesis for increased extracellular L-glutamate concentration in neuroinflammation, *J. Neuroinflammation* 9 (2012) 275.
- [364] Y. Ma, W. Cao, L. Wang, J. Jiang, H. Nie, B. Wang, X. Wei, W. Ying, Basal CD38/cyclic ADP-ribose-dependent signaling mediates ATP release and survival of microglia by modulating connexin 43 hemichannels, *Glia* 62 (2014) 943–955.
- [365] K. Dobrenis, H.Y. Chang, M.H. Pina-Benabou, A. Woodroffe, S.C. Lee, R. Rozental, D.C. Spray, E. Scemes, Human and mouse microglia express connexin36, and functional gap junctions are formed between rodent microglia and neurons, *J. Neurosci. Res.* 82 (2005) 306–315.
- [366] S.K. Wasseff, S.S. Scherer, Activated microglia do not form functional gap junctions in vivo, *J. Neuroimmunol.* 269 (2014) 90–93.
- [367] W. Meme, C.F. Calvo, N. Froger, P. Ezan, E. Amigou, A. Koulakoff, C. Giaume, Proinflammatory cytokines released from microglia inhibit gap junctions in astrocytes: potentiation by beta-amyloid, *FASEB J.* 20 (2006) 494–496.
- [368] J. Xiong, M. Burkovetskaya, N. Karpuk, T. Kielian, IL-1RI (interleukin-1 receptor type I) signalling is essential for host defence and hemichannel activity during acute central nervous system bacterial infection, *ASN Neuro* 4 (2012).
- [369] F.F. Zhang, N. Morioka, K. Nakashima-Hisaoka, Y. Nakata, Spinal astrocytes stimulated by tumor necrosis factor- $\alpha$  and/or interferon- $\gamma$  attenuate connexin 43-gap junction via c-jun terminal kinase activity, *J. Neurosci. Res.* 91 (2013) 745–756.
- [370] G.R. John, E. Scemes, S.O. Suadcani, J.S. Liu, P.C. Charles, S.C. Lee, D.C. Spray, C.F. Brosnan, IL-1 $\beta$  differentially regulates calcium wave propagation between primary human fetal astrocytes via pathways involving P2 receptors and gap junction channels, *Proc. Natl. Acad. Sci. U. S. A.* 96 (1999) 11613–11618.
- [371] J.A. Orellana, T.D. Montero, R. von Bernhardt, Astrocytes inhibit nitric oxide-dependent Ca(2+) dynamics in activated microglia: involvement of ATP released via pannexin 1 channels, *Glia* 61 (2013) 2023–2037.
- [372] E.A. Eugenin, D. Eckardt, M. Theis, K. Willecke, M.V. Bennett, J.C. Saez, Microglia at brain stab wounds express connexin 43 and in vitro form functional gap junctions after treatment with interferon- $\gamma$  and tumor necrosis factor- $\alpha$ , *Proc. Natl. Acad. Sci. U. S. A.* 98 (2001) 4190–4195.
- [373] P.J. Saez, K.F. Shoji, M.A. Retamal, P.A. Harcha, G. Ramirez, J.X. Jiang, R. von Bernhardt, J.C. Saez, ATP is required and advances cytokine-induced gap junction formation in microglia in vitro, *Mediat. Inflamm.* 2013 (2013) 216402.
- [374] D. Davalos, J. Grutzendler, G. Yang, J.V. Kim, Y. Zuo, S. Jung, D.R. Littman, M.L. Dustin, W.B. Gan, ATP mediates rapid microglial response to local brain injury in vivo, *Nat. Neurosci.* 8 (2005) 752–758.
- [375] C. Verderio, M. Matteoli, ATP mediates calcium signaling between astrocytes and microglial cells: modulation by IFN- $\gamma$ , *J. Immunol.* 166 (2001) 6383–6391.
- [376] J.V. Kim, M.L. Dustin, Innate response to focal necrotic injury inside the blood–brain barrier, *J. Immunol.* 177 (2006) 5269–5277.
- [377] F.B. Chekeni, M.R. Elliott, J.K. Sandilos, S.F. Walk, J.M. Kinchen, E.R. Lazarowski, A.J. Armstrong, S. Penuela, D.W. Laird, G.S. Salvesen, B.E. Isakson, D.A. Bayliss, K.S. Ravichandran, Pannexin 1 channels mediate 'find-me' signal release and membrane permeability during apoptosis, *Nature* 467 (2010) 863–867.
- [378] Y. Qu, S. Misaghi, K. Newton, L.L. Gilmour, S. Louie, J.E. Cupp, G.R. Dubyak, D. Hackos, V.M. Dixit, Pannexin-1 is required for ATP release during apoptosis but not for inflammasome activation, *J. Immunol.* 186 (2011) 6553–6561.
- [379] W.R. Silverman, J.P. de Rivero Vaccari, S. Locovei, F. Qiu, S.K. Carlsson, E. Scemes, R. W. Keane, G. Dahl, The pannexin 1 channel activates the inflammasome in neurons and astrocytes, *J. Biol. Chem.* 284 (2009) 18143–18151.
- [380] M.F. Santiago, J. Veliskova, N.K. Patel, S.E. Lutz, D. Caille, A. Charollais, P. Meda, E. Scemes, Targeting pannexin1 improves seizure outcome, *PLoS One* 6 (2011) e25178.
- [381] S.E. Samuels, J.B. Lipitz, J. Wang, G. Dahl, K.J. Muller, Arachidonic acid closes innexin/pannexin channels and thereby inhibits microglia cell movement to a nerve injury, *Dev. Neurobiol.* 73 (2013) 621–631.
- [382] J. Xia, J.C. Lim, W. Lu, J.M. Beckel, E.J. Macarak, A.M. Laties, C.H. Mitchell, Neurons respond directly to mechanical deformation with pannexin-mediated ATP release and autostimulation of P2X7 receptors, *J. Physiol.* 590 (2012) 2285–2304.
- [383] A. Nimmerjahn, F. Kirchhoff, F. Helmchen, Resting microglial cells are highly dynamic surveillants of brain parenchyma in vivo, *Science* 308 (2005) 1314–1318.
- [384] Y. Li, X.F. Du, J.L. Du, Resting microglia respond to and regulate neuronal activity in vivo, *Commun. Integr. Biol.* 6 (2013) e24493.
- [385] E.M. Blanc, A.J. Bruce-Keller, M.P. Mattson, Astrocytic gap junctional communication decreases neuronal vulnerability to oxidative stress-induced disruption of Ca<sup>2+</sup> homeostasis and cell death, *J. Neurochem.* 70 (1998) 958–970.
- [386] T. Nakase, S. Fushiki, C.C. Naus, Astrocytic gap junctions composed of connexin 43 reduce apoptotic neuronal damage in cerebral ischemia, *Stroke* 34 (2003) 1987–1993.
- [387] M.V. Frantseva, L. Kokarovtseva, C.G. Naus, P.L. Carlen, D. MacFabe, J.L. Perez Velazquez, Specific gap junctions enhance the neuronal vulnerability to brain traumatic injury, *J. Neurosci.* 22 (2002) 644–653.
- [388] J.J. Yoon, C.R. Green, S.J. O'Carroll, L.F. Nicholson, Dose-dependent protective effect of connexin43 mimetic peptide against neurodegeneration in an ex vivo model of epileptiform lesion, *Epilepsy Res.* 92 (2010) 153–162.
- [389] T. Unger, S. Bette, J. Zhang, M. Theis, J. Engele, Connexin-deficiency affects expression levels of glial glutamate transporters within the cerebrum, *Neurosci. Lett.* 506 (2012) 12–16.
- [390] J. Iyyathurai, C. D'Hondt, N. Wang, M. De Bock, B. Himpens, M.A. Retamal, J. Stehberg, L. Leybaert, G. Bultynck, Peptides and peptide-derived molecules targeting the intracellular domains of Cx43: Gap junctions versus hemichannels, *Neuropharmacology* 75 (2013) 491–505.
- [391] N. Wang, M. De Bock, E. Decrock, M. Bol, A. Gadicherla, G. Bultynck, L. Leybaert, Connexin targeting peptides as inhibitors of voltage- and intracellular Ca-triggered Cx43 hemichannel opening, *Neuropharmacology* 75 (2013) 506–516.



- [392] W.H. Evans, G. Bultynck, L. Leybaert, Manipulating connexin communication channels: use of peptidomimetics and the translational outputs, *J. Membr. Biol.* 245 (2012) 437–449.
- [393] N. Wang, M. De Bock, G. Antoons, A.K. Gadicherla, M. Bol, E. Decrock, W.H. Evans, K. R. Sipido, F.F. Bukauskas, L. Leybaert, Connexin mimetic peptides inhibit Cx43 hemichannel opening triggered by voltage and intracellular  $\text{Ca}^{2+}$  elevation, *Basic Res. Cardiol.* 107 (2012) 304.
- [394] M. De Bock, M. Culot, N. Wang, M. Bol, E. Decrock, E. De Vuyst, A. da Costa, I. Dauwe, M. Vinken, A.M. Simon, V. Rogiers, G. De Ley, W.H. Evans, G. Bultynck, G. Dupont, R. Cecchelli, L. Leybaert, Connexin channels provide a target to manipulate brain endothelial calcium dynamics and blood–brain barrier permeability, *J. Cereb. Blood Flow Metab.* 31 (2011) 1942–1957.
- [395] P.E. Martin, C. Wall, T.M. Griffith, Effects of connexin-mimetic peptides on gap junction functionality and connexin expression in cultured vascular cells, *Br. J. Pharmacol.* 144 (2005) 617–627.
- [396] N. Karpuk, M. Burkovetskaya, T. Fritz, A. Angle, T. Kielian, Neuroinflammation leads to region-dependent alterations in astrocyte gap junction communication and hemichannel activity, *J. Neurosci.* 31 (2011) 414–425.
- [397] R. Ponsaerts, E. De Vuyst, M. Retamal, C. D'Hondt, D. Vermeire, N. Wang, H. De Smedt, P. Zimmermann, B. Himpens, J. Vereecke, L. Leybaert, G. Bultynck, Intramolecular loop/tail interactions are essential for connexin 43-hemichannel activity, *FASEB J.* 24 (2010) 4378–4395.
- [398] N. Wang, E. De Vuyst, R. Ponsaerts, K. Boengler, N. Palacios-Prado, J. Wauman, C.P. Lai, M. De Bock, E. Decrock, M. Bol, M. Vinken, V. Rogiers, J. Tavernier, W.H. Evans, C.C. Naus, F.F. Bukauskas, K.R. Sipido, G. Heusch, R. Schulz, G. Bultynck, L. Leybaert, Selective inhibition of Cx43 hemichannels by Gap19 and its impact on myocardial ischemia/reperfusion injury, *Basic Res. Cardiol.* 108 (2012) 309.
- [399] J.O. Davidson, C.R. Green, L.F. Nicholson, S.J. O'Carroll, M. Fraser, L. Bennet, A.J. Gunn, Connexin hemichannel blockade improves outcomes in a model of fetal ischemia, *Ann. Neurol.* 71 (2012) 121–132.
- [400] J.A. Orellana, D.E. Hernandez, P. Ezan, V. Velarde, M.V. Bennett, C. Giaume, J.C. Saez, Hypoxia in high glucose followed by reoxygenation in normal glucose reduces the viability of cortical astrocytes through increased permeability of connexin 43 hemichannels, *Glia* 58 (2010) 329–343.
- [401] M. Samoilova, K. Wentlandt, Y. Adamchik, A.A. Velumian, P.L. Carlen, Connexin 43 mimetic peptides inhibit spontaneous epileptiform activity in organotypic hippocampal slice cultures, *Exp. Neurol.* 210 (2008) 762–775.
- [402] J.A. Orellana, K.F. Shoji, V. Abudara, P. Ezan, E. Amigou, P.J. Saez, J.X. Jiang, C.C. Naus, J.C. Saez, C. Giaume, Amyloid beta-induced death in neurons involves glial and neuronal hemichannels, *J. Neurosci.* 31 (2011) 4962–4977.
- [403] Y.S. Chen, I. Toth, H.V. Danesh-Meyer, C.R. Green, I.D. Rupenthal, Cytotoxicity and vitreous stability of chemically modified connexin43 mimetic peptides for the treatment of optic neuropathy, *J. Pharm. Sci.* 102 (2013) 2322–2331.
- [404] S.J. O'Carroll, M. Alkadhi, L.F. Nicholson, C.R. Green, Connexin 43 mimetic peptides reduce swelling, astrogliosis, and neuronal cell death after spinal cord injury, *Cell Commun. Adhes.* 15 (2008) 27–42.
- [405] P.L. Carlen, Curious and contradictory roles of glial connexins and pannexins in epilepsy, *Brain Res.* 1487 (2012) 54–60.
- [406] A. Koulakoff, X. Mei, J.A. Orellana, J.C. Saez, C. Giaume, Glial connexin expression and function in the context of Alzheimer's disease, *Biochim. Biophys. Acta* 1818 (2012) 2048–2057.
- [407] J.A. Orellana, R. von Bernhardt, C. Giaume, J.C. Saez, Glial hemichannels and their involvement in aging and neurodegenerative diseases, *Rev. Neurosci.* 23 (2012) 163–177.
- [408] J.O. Davidson, C.R. Green, L. Bennet, L.F. Nicholson, H. Danesh-Meyer, S.J. O'Carroll, A.J. Gunn, A key role for connexin hemichannels in spreading ischemic brain injury, *Curr. Drug Targets* 14 (2013) 36–46.
- [409] R.A. Quintanilla, J.A. Orellana, R. von Bernhardt, Understanding risk factors for Alzheimer's disease: interplay of neuroinflammation, connexin-based communication and oxidative stress, *Arch. Med. Res.* 43 (2013) 632–644.
- [410] T. Nakase, T. Maeda, Y. Yoshida, K. Nagata, Ischemia alters the expression of connexins in the aged human brain, *J. Biomed. Biotechnol.* 2009 (2009) 147946.
- [411] M.A. Moskowitz, E.H. Lo, C. Iadecola, The science of stroke: mechanisms in search of treatments, *Neuron* 67 (2010) 181–198.
- [412] R.C. Turner, S.C. Dodson, C.L. Rosen, J.D. Huber, The science of cerebral ischemia and the quest for neuroprotection: navigating past failure to future success, *J. Neurosurg.* 118 (2013) 1072–1085.
- [413] J. Astrup, B.K. Siesjö, L. Symon, Thresholds in cerebral ischemia—the ischemic penumbra, *Stroke* 12 (1981) 723–725.
- [414] J. Astrup, L. Symon, N.M. Branstetter, N.A. Lassen, Cortical evoked potential and extracellular  $\text{K}^{+}$  and  $\text{H}^{+}$  at critical levels of brain ischemia, *Stroke* 8 (1977) 51–57.
- [415] E.H. Lo, A new penumbra: transitioning from injury into repair after stroke, *Nat. Med.* 14 (2008) 497–500.
- [416] G. Barreto, R.E. White, Y. Ouyang, L. Xu, R.G. Giffard, Astrocytes: targets for neuroprotection in stroke, *Cent. Nerv. Syst. Agents Med. Chem.* 11 (2011) 164–173.
- [417] Y. Zhao, D.A. Rempe, Targeting astrocytes for stroke therapy, *Neurotherapeutics* 7 (2010) 439–451.
- [418] T. Takano, N. Oberheim, M.L. Cotrina, M. Nedergaard, Astrocytes and ischemic injury, *Stroke* 40 (2009) S8–S12.
- [419] A.M. Brown, Brain glycogen re-awakened, *J. Neurochem.* 89 (2004) 537–552.
- [420] K. Kazama, G. Wang, K. Frys, J. Anrather, C. Iadecola, Angiotensin II attenuates functional hyperemia in the mouse somatosensory cortex, *Am. J. Physiol. Heart Circ. Physiol.* 285 (2003) H1890–H1899.
- [421] K. Niwa, L. Younkin, C. Ebeling, S.K. Turner, D. Westaway, S. Younkin, K.H. Ashe, G. A. Carlson, C. Iadecola, Abeta 1–40-related reduction in functional hyperemia in mouse neocortex during somatosensory activation, *Proc. Natl. Acad. Sci. U. S. A.* 97 (2000) 9735–9740.
- [422] B.V. Zlokovic, The blood–brain barrier in health and chronic neurodegenerative disorders, *Neuron* 57 (2008) 178–201.
- [423] S.S. Chew, C.S. Johnson, C.R. Green, H.V. Danesh-Meyer, Role of connexin43 in central nervous system injury, *Exp. Neurol.* 225 (2010) 250–261.
- [424] P. Bargiotas, H. Monyer, M. Schwanninger, Hemichannels in cerebral ischemia, *Curr. Mol. Med.* 9 (2009) 186–194.
- [425] M. Sahores, A. Mendoza-Naranjo, Gap junctions as therapeutic targets in brain injury following hypoxia–ischemia, *Recent Pat. CNS Drug Discov.* 3 (2008) 209–215.
- [426] J.E. Contreras, H.A. Sanchez, L.P. Veliz, F.F. Bukauskas, M.V. Bennett, J.C. Saez, Role of connexin-based gap junction channels and hemichannels in ischemia-induced cell death in nervous tissue, *Brain Res. Brain Res. Rev.* 47 (2004) 290–303.
- [427] Y. Wang, J.V. Denisova, K.S. Kang, J.D. Fontes, B.T. Zhu, A.B. Belousov, Neuronal gap junctions are required for NMDA receptor-mediated excitotoxicity: implications in ischemic stroke, *J. Neurophysiol.* 104 (2010) 3551–3556.
- [428] M.A. Retamal, K.A. Schalper, K.F. Shoji, J.A. Orellana, M.V. Bennett, J.C. Saez, Possible involvement of different connexin43 domains in plasma membrane permeabilization induced by ischemia–reperfusion, *J. Membr. Biol.* 218 (2007) 49–63.
- [429] X. Wang, A. Ma, W. Zhu, L. Zhu, Y. Zhao, J. Xi, X. Zhang, B. Zhao, The role of connexin 43 and hemichannels correlated with the astrocytic death following ischemia/reperfusion insult, *Cell. Mol. Neurobiol.* 33 (2013) 401–410.
- [430] M.L. Cotrina, J. Kang, J.H. Lin, E. Bueno, T.W. Hansen, L. He, Y. Liu, M. Nedergaard, Astrocytic gap junctions remain open during ischemic conditions, *J. Neurosci.* 18 (1998) 2520–2537.
- [431] M. Nedergaard, Spreading depression as a contributor to ischemic brain damage, *Adv. Neurol.* 71 (1996) 75–83 (discussion 83–74).
- [432] M. Nedergaard, A.J. Cooper, S.A. Goldman, Gap junctions are required for the propagation of spreading depression, *J. Neurobiol.* 28 (1995) 433–444.
- [433] A. Rawanduzay, A. Hansen, T.W. Hansen, M. Nedergaard, Effective reduction of infarct volume by gap junction blockade in a rodent model of stroke, *J. Neurosurg.* 87 (1997) 916–920.
- [434] T.A. Basarsky, S.N. Duffy, R.D. Andrew, B.A. MacVicar, Imaging spreading depression and associated intracellular calcium waves in brain slices, *J. Neurosci.* 18 (1998) 7189–7199.
- [435] P.E. Kunkler, R.P. Kraig, Calcium waves precede electrophysiological changes of spreading depression in hippocampal organ cultures, *J. Neurosci.* 18 (1998) 3416–3425.
- [436] M. Tanaka, K. Kawahara, T. Kosugi, T. Yamada, T. Mioka, Changes in the spontaneous calcium oscillations for the development of the preconditioning-induced ischemic tolerance in neuron/astrocyte co-culture, *Neurochem. Res.* 32 (2007) 988–1001.
- [437] L. Tao, A.L. Harris, 2-aminoethoxydiphenyl borate directly inhibits channels composed of connexin26 and/or connexin32, *Mol. Pharmacol.* 71 (2007) 570–579.
- [438] M.D. Bootman, T.J. Collins, L. Mackenzie, H.L. Roderick, M.J. Berridge, C.M. Peppiatt, 2-aminoethoxydiphenyl borate (2-APB) is a reliable blocker of store-operated  $\text{Ca}^{2+}$  entry but an inconsistent inhibitor of  $\text{InsP}_3$ -induced  $\text{Ca}^{2+}$  release, *FASEB J.* 16 (2002) 1145–1150.
- [439] Y. Takatsuru, K. Eto, R. Kaneko, H. Masuda, N. Shimokawa, N. Koibuchi, J. Nabekura, Critical role of the astrocyte for functional remodeling in contralateral hemisphere of somatosensory cortex after stroke, *J. Neurosci.* 33 (2013) 4683–4692.
- [440] M.J. Berridge, Calcium hypothesis of Alzheimer's disease, *Pflügers Arch.* 459 (2010) 441–449.
- [441] S. Chakraborty, G.E. Stutzmann, Early calcium dysregulation in Alzheimer's disease: setting the stage for synaptic dysfunction, *Sci. China Life Sci.* 54 (2011) 752–762.
- [442] K. Furukawa, Y. Wang, P.J. Yao, W. Fu, M.P. Mattson, Y. Itoyama, H. Onodera, I. D'Souza, P.H. Poorkaj, T.D. Bird, G.D. Schellenberg, Alteration in calcium channel properties is responsible for the neurotoxic action of a familial frontotemporal dementia tau mutation, *J. Neurochem.* 87 (2003) 427–436.
- [443] H. Tu, O. Nelson, A. Bezprozvanny, Z. Wang, S.F. Lee, Y.H. Hao, L. Serneels, B. De Strooper, G. Yu, I. Bezprozvanny, Presenilins form  $\text{ER Ca}^{2+}$  leak channels, a function disrupted by familial Alzheimer's disease-linked mutations, *Cell* 126 (2006) 981–993.
- [444] V. Rybalchenko, S.Y. Hwang, N. Rybalchenko, P. Koulen, The cytosolic N-terminus of presenilin-1 potentiates mouse ryanodine receptor single channel activity, *Int. J. Biochem. Cell Biol.* 40 (2008) 84–97.
- [445] K.N. Green, A. Demuro, Y. Akbari, B.D. Hitt, I.F. Smith, I. Parker, F.M. LaFerla, SERCA pump activity is physiologically regulated by presenilin and regulates amyloid beta production, *J. Cell Biol.* 181 (2008) 1107–1116.
- [446] M.P. Mattson, B. Cheng, D. Davis, K. Bryant, I. Lieberburg, R.E. Rydel, beta-Amyloid peptides destabilize calcium homeostasis and render human cortical neurons vulnerable to excitotoxicity, *J. Neurosci.* 12 (1992) 376–389.
- [447] S.M. Jin, H.J. Cho, Y.W. Kim, J.Y. Hwang, I. Mook-Jung, Abeta-induced  $\text{Ca}^{2+}$  influx regulates astrocytic BACE1 expression via calcineurin/NFAT4 signals, *Biochem. Biophys. Res. Commun.* 425 (2012) 649–655.
- [448] R. Bhatia, H. Lin, R. Lal, Fresh and globular amyloid beta protein (1–42) induces rapid cellular degeneration: evidence for AbetaP channel-mediated cellular toxicity, *FASEB J.* 14 (2000) 1233–1243.
- [449] T. Thomas, G. Thomas, C. McLendon, T. Sutton, M. Mullan, beta-Amyloid-mediated vasoactivity and vascular endothelial damage, *Nature* 380 (1996) 168–171.
- [450] A. Verkhratsky, M. Olabarria, H.N. Noristani, C.Y. Yeh, J.J. Rodriguez, Astrocytes in Alzheimer's disease, *Neurotherapeutics* 7 (2010) 399–412.
- [451] M. Olabarria, H.N. Noristani, A. Verkhratsky, J.J. Rodriguez, Concomitant astroglial atrophy and astrogliosis in a triple transgenic animal model of Alzheimer's disease, *Glia* 58 (2010) 831–838.

- [452] A. Serrano-Pozo, M.L. Mielke, T. Gomez-Isla, R.A. Betensky, J.H. Growdon, M.P. Frosch, B.T. Hyman, Reactive glia not only associates with plaques but also parallels tangles in Alzheimer's disease, *Am. J. Pathol.* 179 (2011) 1373–1384.
- [453] R.G. Nagele, J. Wegiel, V. Venkataraman, H. Imaki, K.C. Wang, J. Wegiel, Contribution of glial cells to the development of amyloid plaques in Alzheimer's disease, *Neurobiol. Aging* 25 (2004) 663–674.
- [454] A.W. Kraft, X. Hu, H. Yoon, P. Yan, Q. Xiao, Y. Wang, S.C. Gil, J. Brown, U. Wilhelmsson, J.L. Restivo, J.R. Cirrito, D.M. Holtzman, J. Kim, M. Pekny, J.M. Lee, Attenuating astrocyte activation accelerates plaque pathogenesis in APP/PS1 mice, *FASEB J.* 27 (2013) 187–198.
- [455] T. Tomida, K. Hirose, A. Takizawa, F. Shibasaki, M. Iino, NFAT functions as a working memory of  $\text{Ca}^{2+}$  signals in decoding  $\text{Ca}^{2+}$  oscillation, *EMBO J.* 22 (2003) 3825–3832.
- [456] P. Tompa, R. Toth-Boconadi, P. Friedrich, Frequency decoding of fast calcium oscillations by calpain, *Cell Calcium* 29 (2001) 161–170.
- [457] D. Lim, A. Iyer, V. Ronco, A.A. Grolla, P.L. Canonico, E. Aronica, A.A. Genazzani, Amyloid beta deregulates astroglial mGluR5-mediated calcium signaling via calcineurin and NF- $\kappa$ B, *Glia* 61 (2013) 1134–1145.
- [458] J.I. Nagy, W. Li, E.L. Hertzberg, C.A. Marotta, Elevated connexin43 immunoreactivity at sites of amyloid plaques in Alzheimer's disease, *Brain Res.* 717 (1996) 173–178.
- [459] N.F. Cruz, K.K. Ball, G.A. Dienel, Astrocytic gap junctional communication is reduced in amyloid-beta-treated cultured astrocytes, but not in Alzheimer's disease transgenic mice, *ASN Neuro* 2 (2010) e00041.
- [460] J.T. Weber, Altered calcium signaling following traumatic brain injury, *Front. Pharmacol.* 3 (2012) 60.
- [461] M.W. Greve, B.J. Zink, Pathophysiology of traumatic brain injury, *Mt Sinai J. Med.* 76 (2009) 97–104.
- [462] Z. Yong-Ming, L. Jia-Chuan, Y. Yan-Yan, S. Wen-Jiang, T. Hong, L. Bing-Cang, Z. Liang-Chao, Effective protection of rabbits' explosive brain injury through blocking gap junction communication, *Afr. Health Sci.* 12 (2013) 552–556.
- [463] Y. Wang, J.H. Song, J.V. Denisova, W.M. Park, J.D. Fontes, A.B. Belousov, Neuronal gap junction coupling is regulated by glutamate and plays critical role in cell death during neuronal injury, *J. Neurosci.* 32 (2012) 713–725.
- [464] A.B. Belousov, Y. Wang, J.H. Song, J.V. Denisova, N.E. Berman, J.D. Fontes, Neuronal gap junctions play a role in the secondary neuronal death following controlled cortical impact, *Neurosci. Lett.* 524 (2012) 16–19.
- [465] A. Shunmugavel, M. Khan, P.C. Chou, I. Singh, Spinal cord injury induced arrest in estrous cycle of rats is ameliorated by S-nitrosoglutathione: novel therapeutic agent to treat amenorrhea, *J. Sex. Med.* 9 (2012) 148–158.
- [466] A. Ohsumi, H. Nawashiro, N. Otani, H. Ooigawa, T. Toyooka, K. Shima, Temporal and spatial profile of phosphorylated connexin43 after traumatic brain injury in rats, *J. Neurotrauma* 27 (2010) 1255–1263.
- [467] L.Q. Sun, J.L. Gao, C.M. Cui, Y. Cui, X.B. Jing, M.M. Zhao, Y.C. Wang, Y.X. Tian, K.J. Wang, J.Z. Cui, Astrocytic p-connexin 43 regulates neuronal autophagy in the hippocampus following traumatic brain injury in rats, *Mol. Med. Rep.* 9 (2014) 77–82.
- [468] Y. Moon, S.Y. Choi, K. Kim, H. Kim, W. Sun, Expression of connexin29 and 32 in the penumbra region after traumatic brain injury of mice, *Neuroreport* 21 (2010) 1135–1139.
- [469] C.L. Floyd, B.A. Rzigalinski, J.T. Weber, H.A. Sitterding, K.A. Willoughby, E.F. Ellis, Traumatic injury of cultured astrocytes alters inositol (1,4,5)-trisphosphate-mediated signaling, *Glia* 33 (2001) 12–23.
- [470] C.W. Taylor, O. Dellis, Plasma membrane IP3 receptors, *Biochem. Soc. Trans.* 34 (2006) 910–912.
- [471] U. Heinemann, J. Louvel, Changes in  $[\text{Ca}^{2+}]_o$  and  $[\text{K}^{+}]_o$  during repetitive electrical stimulation and during pentetrazol induced seizure activity in the sensorimotor cortex of cats, *Pflugers Arch.* 398 (1983) 310–317.
- [472] Y. Yaari, A. Konnerth, U. Heinemann, Spontaneous epileptiform activity of CA1 hippocampal neurons in low extracellular calcium solutions, *Exp. Brain Res.* 51 (1983) 153–156.
- [473] S.H. Lee, S. Magge, D.D. Spencer, H. Sontheimer, A.H. Cornell-Bell, Human epileptic astrocytes exhibit increased gap junction coupling, *Glia* 15 (1995) 195–202.
- [474] M.H. De Pina-Benabou, M. Srinivas, D.C. Spray, E. Scemes, Calmodulin kinase pathway mediates the  $\text{K}^{+}$ -induced increase in Gap junctional communication between mouse spinal cord astrocytes, *J. Neurosci.* 21 (2001) 6635–6643.
- [475] R. Kohling, S.J. Gladwell, E. Bracci, M. Vreugdenhil, J.G. Jefferys, Prolonged epileptiform bursting induced by 0-Mg(2+) in rat hippocampal slices depends on gap junctional coupling, *Neuroscience* 105 (2001) 579–587.
- [476] N. Kang, J. Xu, Q. Xu, M. Nedergaard, J. Kang, Astrocytic glutamate release-induced transient depolarization and epileptiform discharges in hippocampal CA1 pyramidal neurons, *J. Neurophysiol.* 94 (2005) 4121–4130.
- [477] T. Fellin, M. Gomez-Gonzalo, S. Gobbo, G. Carmignoto, P.G. Haydon, Astrocytic glutamate is not necessary for the generation of epileptiform neuronal activity in hippocampal slices, *J. Neurosci.* 26 (2006) 9312–9322.